

NR3C1 Antibody (C-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP4867b

Specification

NR3C1 Antibody (C-term) - Product Information

Application	IF, WB, IHC-P,E
Primary Accession	P04150
Other Accession	P04150-2
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit Ig
Calculated MW	85659
Antigen Region	713-742

NR3C1 Antibody (C-term) - Additional Information

Gene ID 2908

Other Names

Glucocorticoid receptor, GR, Nuclear receptor subfamily 3 group C member 1, NR3C1, GRL

Target/Specificity

This NR3C1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 713-742 amino acids from the C-terminal region of human NR3C1.

Dilution

IF~~1:10~50
WB~~1:1000
IHC-P~~1:50~100


Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.


Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions



Confocal immunofluorescent analysis of NR3C1 Antibody (C-term)(Cat#AP4867b) with HeLa cell followed by Alexa Fluor 488-conjugated goat anti-rabbit IgG (green). Actin filaments have been labeled with Alexa Fluor 555 phalloidin (red). DAPI was used to stain the cell nuclear (blue).



Western blot analysis of NR3C1 Antibody (C-term) (Cat. #AP4867b) in A549 cell line lysates (35ug/lane). NR3C1 (arrow) was detected using the purified Pab.

NR3C1 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

NR3C1 Antibody (C-term) - Protein Information

Name NR3C1


Synonyms GRL

Function

Receptor for glucocorticoids (GC) (PubMed:27120390). Has a dual mode of action: as a transcription factor that binds to glucocorticoid response elements (GRE), both for nuclear and mitochondrial DNA, and as a modulator of other transcription factors. Affects inflammatory responses, cellular proliferation and differentiation in target tissues. Involved in chromatin remodeling (PubMed:9590696). Plays a role in rapid mRNA degradation by binding to the 5' UTR of target mRNAs and interacting with PNRC2 in a ligand-dependent manner which recruits the RNA helicase UPF1 and the mRNA- decapping enzyme DCP1A, leading to RNA decay (PubMed:25775514). Could act as a coactivator for STAT5-dependent transcription upon growth hormone (GH) stimulation and could reveal an essential role of hepatic GR in the control of body growth (By similarity).

Cellular Location

[Isoform Alpha]: Cytoplasm. Nucleus. Mitochondrion. Cytoplasm, cytoskeleton, spindle. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome. Note=After ligand activation, translocates from the cytoplasm to the nucleus. In the presence of NR1D1 shows a time-dependent subcellular localization, localizing to the cytoplasm at ZT8 and to the nucleus at ZT20 (By similarity). Lacks this diurnal pattern of localization in the absence of NR1D1, localizing to both nucleus and the cytoplasm at ZT8 and ZT20 (By similarity).



NR3C1 Antibody (C-term) (Cat. #AP4867B) immunohistochemistry analysis in formalin fixed and paraffin embedded human kidney tissue followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of NR3C1 Antibody (C-term) for immunohistochemistry. Clinical relevance has not been evaluated.

NR3C1 Antibody (C-term) - Background

NR3C1 is a receptor for glucocorticoids that can act as both a transcription factor and as a regulator of other transcription factors. This protein can also be found in heteromeric cytoplasmic complexes along with heat shock factors and immunophilins. The protein is typically found in the cytoplasm until it binds a ligand, which induces transport into the nucleus. Mutations in this gene are a cause of glucocorticoid resistance, or cortisol, resistance.

NR3C1 Antibody (C-term) - References

- Geelhoed, M.J., et al. BMC Med. Genet. 11, 39 (2010)
- Szilagyi, K., et al. Neuro Endocrinol. Lett. 30(5):629-636(2009)
- Tian, S., et al. Biochem. J. 367 (PT 3), 907-911 (2002)
- Itoh, M., et al. Mol. Endocrinol. 16(10):2382-2392(2002)
- Wang, Z., et al. J. Biol. Chem. 277(29):26573-26580(2002)

{ECO:0000250|UniProtKB:P06537,
ECO:0000269|PubMed:18838540,
ECO:0000269|PubMed:27120390,
ECO:0000269|PubMed:8621628} [Isoform
Alpha-B]: Nucleus. Cytoplasm Note=After
ligand activation, translocates from the
cytoplasm to the nucleus.

Tissue Location

Widely expressed including bone, stomach,
lung, liver, colon, breast, ovary, pancreas
and kidney (PubMed:25847991). In the
heart, detected in left and right atria, left
and right ventricles, aorta, apex,
intraventricular septum, and
atrioventricular node as well as whole adult
and fetal heart (PubMed:10902803)
[Isoform Alpha-2]: Widely expressed.

NR3C1 Antibody (C-term) - Protocols

Provided below are standard protocols that you
may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)