

CASP3(Asp175) Antibody
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP4985D

Specification

CASP3(Asp175) Antibody - Product Information

Application	WB, IHC-P, FC,E
Primary Accession	P42574
Other Accession	NP_004337
Reactivity	Human, Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit Ig
Antigen Region	149-179

CASP3(Asp175) Antibody - Additional Information

Gene ID 836

Other Names

Caspase-3, CASP-3, Apopain, Cysteine protease CPP32, CPP-32, Protein Yama, SREBP cleavage activity 1, SCA-1, Caspase-3 subunit p17, Caspase-3 subunit p12, CASP3, CPP32

Target/Specificity

This CASP3(Asp175) antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 149-179 amino acids from human CASP3(Asp175).

Dilution


WB~~1:1000-1:2000
IHC-P~~1:50~100
FC~~1:10~50

Format


Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.



Western blot analysis of CASP3 (Asp175) Antibody (Cat. #AP4985d) in CEM cell line lysates (35ug/lane). CASP3 (arrow) was detected using the purified Pab.



Western blot analysis of CASP3 (arrow) using rabbit polyclonal CASP3 Antibody (Cat. #AP4985d). 293 cell lysates (2 ug/lane) either nontransfected (Lane 1) or transiently transfected (Lane 2) with the CASP3 gene.

Precautions

CASP3(Asp175) Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

CASP3(Asp175) Antibody - Protein Information

Name CASP3

Synonyms CPP32

Function

Involved in the activation cascade of caspases responsible for apoptosis execution (PubMed:7596430). At the onset of apoptosis it proteolytically cleaves poly(ADP-ribose) polymerase (PARP) at a '216- Asp-Gly-217' bond (PubMed:7774019). Cleaves and activates sterol regulatory element binding proteins (SREBPs) between the basic helix-loop-helix leucine zipper domain and the membrane attachment domain. Cleaves and activates caspase-6, -7 and -9 (PubMed:7596430). Involved in the cleavage of huntingtin (PubMed:8696339). Triggers cell adhesion in sympathetic neurons through RET cleavage (PubMed:21357690). Cleaves and inhibits serine/threonine-protein kinase AKT1 in response to oxidative stress (PubMed:23152800).


Cellular Location

Cytoplasm.


Tissue Location

Highly expressed in lung, spleen, heart, liver and kidney. Moderate levels in brain and skeletal muscle, and low in testis. Also found in many cell lines, highest expression in cells of the immune system

CASP3(Asp175) Antibody - Protocols




Anti-CASP3(Asp175) Antibody at 1:1000 dilution + Rat heart whole tissue lysate
Lysates/proteins at 20 µg per lane.
Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution.
Predicted band size : 35 kDa
Blocking/Dilution buffer: 5% NFDM/TBST.




All lanes : Anti-CASP3(Asp175) Antibody at 1:1000-1:2000 dilution
Lane 1: Jurkat whole cell lysate
Lane 2: Rat heart tissue lysate
Lysates/proteins at 20 µg per lane.
Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution.
Predicted band size : 32 kDa
Blocking/Dilution buffer: 5% NFDM/TBST.

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)



CASP3(Asp175) Antibody (Cat. #AP4985d) IHC analysis in formalin fixed and paraffin embedded human brain tissue followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of the CASP3(Asp175) Antibody for immunohistochemistry. Clinical relevance has not been evaluated.



CASP3(Asp175) Antibody (Cat. #AP4985d) flow cytometric analysis of CEM cells (right histogram) compared to a negative control cell (left histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

CASP3(Asp175) Antibody - Background

This gene encodes a protein which is a member of the cysteine-aspartic acid protease (caspase) family. Sequential activation of caspases plays a central role in the execution-phase of cell apoptosis. Caspases exist as inactive proenzymes which undergo proteolytic processing at conserved aspartic residues to produce two subunits, large and

small, that dimerize to form the active enzyme. This protein cleaves and activates caspases 6, 7 and 9, and the protein itself is processed by caspases 8, 9 and 10. It is the predominant caspase involved in the cleavage of amyloid-beta 4A precursor protein, which is associated with neuronal death in Alzheimer's disease. Alternative splicing of this gene results in two transcript variants that encode the same protein.

CASP3(Asp175) Antibody - References

Mei, Y., et al. Mol. Cell 37(5):668-678(2010)
Sohn, E.J., et al. Cancer Res. 70(3):1154-1163(2010) Karamitopoulou, E., et al. Pathology 42(1):37-42(2010) Wang, W., et al. Xi Bao Yu Fen Zi Mian Yi Xue Za Zhi 25(11):1034-1035(2009)