

PYY Antibody (C-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP5103B

Specification

PYY Antibody (C-term) - Product Information

Application IF, WB, IHC-P,

FC,E

Primary Accession
Reactivity
Host
Clonality
Isotype
Calculated MW
Antigen Region

P10082
Human
Rabbit
Polyclonal
Rabbit Ig
11145
67-94

PYY Antibody (C-term) - Additional Information

Gene ID 5697

Other Names

Peptide YY, PYY, PYY-I, PYY

Target/Specificity

This PYY antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 67-94 amino acids from the C-terminal region of human PYY.

Dilution

IF~~1:10~50 WB~~1:1000 IHC-P~~1:10~50 FC~~1:10~50

Format


Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage


Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

PYY Antibody (C-term) is for research use only and not for use in diagnostic or



Confocal immunofluorescent analysis of PYY Antibody (C-term)(Cat. #AP5103b) with MCF-7 cell followed by Alexa Fluor[]?488-conjugated goat anti-rabbit IgG (green). DAPI was used to stain the cell nuclear (blue).



Western blot analysis of PYY Antibody (C-term) (Cat. #AP5103b) in MCF-7 cell line lysates (35ug/lane).PYY (arrow) was detected using the purified Pab.



therapeutic procedures.

PYY Antibody (C-term) - Protein Information

Name PYY

Function


This gut peptide inhibits exocrine pancreatic secretion, has a vasoconstrictory action and inhibitis jejunal and colonic mobility.

Cellular Location Secreted.


PYY Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture




PYY Antibody (C-term) (Cat. #AP5103b) IHC analysis in formalin fixed and paraffin embedded colon carcinoma followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of the PYY Antibody (C-term) for immunohistochemistry. Clinical relevance has not been evaluated.




PYY Antibody (C-term) (Cat. #AP5103b) flow cytometric analysis of MCF-7 cells (bottom histogram) compared to a negative control cell (top histogram).FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

PYY Antibody (C-term) - Background

PYY gut peptide inhibits exocrine pancreatic secretion, has a vasoconstrictory action and inhibitis jejunal and colonic mobility.







PYY Antibody (C-term) - References

Ueda, S.Y., et al. J. Endocrinol. 203(3):357-364(2009) Shih, P.A., et al. J. Clin. Endocrinol. Metab. 94(11):4557-4566(2009) Gratacos, M., et al. Am. J. Med. Genet. B Neuropsychiatr. Genet. 150B (6), 808-816 (2009)

PYY Antibody (C-term) - Citations

• G protein βy subunits inhibit TRPM3 ion channels in sensory neurons.