

TUBB2C Antibody (Center)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP5126c

Specification

TUBB2C Antibody (Center) - Product Information

Application	WB, IHC-P,E
Primary Accession	P68371
Other Accession	P09244 , Q91575 , P69897 , Q767L7 , P99024 , P07437 , P69893 , Q2KJD0 , P30883 , Q6P9T8 , P68372 , Q3MHM5 , P09206 , Q24560
Reactivity Predicted	Human Drosophila, Chicken, Bovine, Mouse, Rat, Xenopus, Hamster, Pig
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit Ig
Calculated MW	49831
Antigen Region	99-125

TUBB2C Antibody (Center) - Additional Information

Gene ID 10383

Other Names

Tubulin beta-4B chain, Tubulin beta-2 chain,
Tubulin beta-2C chain, TUBB4B, TUBB2C

Target/Specificity


This TUBB2C antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 99-125 amino acids from the Central region of human TUBB2C.

Dilution


WB~~1:1000
IHC-P~~1:50~100

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A



Western blot analysis of TUBB2C Antibody (Center) (Cat. #AP5126c) in HeLa,MDA-MB435,MDA-MB231,HepG2 cell line lysates (35ug/lane). TUBB2C (arrow) was detected using the purified Pab.



TUBB2C Antibody (Center) (Cat. #AP5126c) IHC analysis in formalin fixed and paraffin embedded human brain tissue followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of the TUBB2C Antibody (Center) for immunohistochemistry. Clinical relevance has not been evaluated.

TUBB2C Antibody (Center) - Background

TUBB2C is the major constituent of

column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

TUBB2C Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

microtubules. It binds two moles of GTP, one at an exchangeable site on the beta chain and one at a non-exchangeable site on the alpha-chain.

TUBB2C Antibody (Center) - References

Xu, W., et al. Mol. Cancer Ther. 8(12):3318-3330(2009)
Chan, C.M., et al. Arch. Pathol. Lab. Med. 132(4):675-683(2008)
Olsen, J.V., et al. Cell 127(3):635-648(2006)

TUBB2C Antibody (Center) - Protein Information

Name TUBB4B

Synonyms TUBB2C

Function

Tubulin is the major constituent of microtubules. It binds two moles of GTP, one at an exchangeable site on the beta chain and one at a non-exchangeable site on the alpha chain.

Cellular Location

Cytoplasm, cytoskeleton.

Tissue Location

Ubiquitous..

TUBB2C Antibody (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)