

CHD3 (C-terminus) Antibody
Purified Mouse Monoclonal Antibody (Mab)
Catalog # AP52685

Specification

CHD3 (C-terminus) Antibody - Product Information

Application	IP, WB, ICC
Primary Accession	Q12873
Reactivity	Human, Mouse
Host	Mouse
Clonality	Monoclonal
Isotype	IgG1
Calculated MW	260 KDa

CHD3 (C-terminus) Antibody - Additional Information

Gene ID 1107

Other Names

226;2600010P09Rik;AF020312;ATP dependent helicase CHD3;ATP-dependent helicase Chd3;CHD-3;Chd3;CHD3_HUMAN;Chd7;Chromodomain-helicase-DNA-binding protein 3;hZFH;MGC40857;Mi 2 autoantigen 240 kDa protein;Mi 2a;Mi-2 autoantigen 240 kDa protein;Mi2 ALPHA;Mi2-alpha;Prp7;Prp9 1;ZFH;Zinc finger helicase;zinc-finger helicase (Snf2-like).

Dilution

IP~~1:500
WB~~1:1000
ICC~~1:400

Format

ascites containing 0.02% NaN₃ & 50% glycerol

Storage


Store at -20 °C.Stable for 12 months from date of receipt

CHD3 (C-terminus) Antibody - Protein Information


Name CHD3

Function

Component of the histone deacetylase



Immunoprecipitation analysis of HeLa cell lysates using CHD3 mouse mAb.



Western blot detection of CHD3 (C-terminus) in NTERA2,3T3,C6,K562 and HeLa cell lysates using CHD3 (C-terminus) mouse mAb (1:1000 diluted).Predicted band size: 260KDa.Observed band size: 260KDa.


NuRD complex which participates in the remodeling of chromatin by deacetylating histones (PubMed:9804427, PubMed:30397230). Involved in transcriptional repression as part of the NuRD complex (PubMed:27068747). Required for anchoring centrosomal pericentrin in both interphase and mitosis, for spindle organization and centrosome integrity (PubMed:17626165).

Cellular Location

Nucleus, PML body. Nucleus. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome. Note=Associates with centrosomes in interphase and mitosis.

Tissue Location

Widely expressed..



Immunocytochemistry staining of HeLa cells fixed with -20°C Methanol and using CHD3 (C-terminus) mouse mAb (dilution 1:400).

CHD3 (C-terminus) Antibody - Background

Component of the histone deacetylase NuRD complex which participates in the remodeling of chromatin by deacetylating histones. Required for anchoring centrosomal pericentrin in both interphase and mitosis, for spindle organization and centrosome integrity.

CHD3 (C-terminus) Antibody - References

Aubry F.,et al.Eur. J. Biochem. 254:558-564(1998).
Zody M.C.,et al.Nature 440:1045-1049(2006).
Mural R.J.,et al.Submitted (SEP-2005) to the EMBL/GenBank/DDBJ databases.
Woodage T.,et al.Proc. Natl. Acad. Sci. U.S.A. 94:11472-11477(1997).
Ge Q.,et al.J. Clin. Invest. 96:1730-1737(1995).

CHD3 (C-terminus) Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)