

Anti-CDK4 Antibody
Mouse Monoclonal Antibody
Catalog # AP53463

Specification

Anti-CDK4 Antibody - Product Information

Application	WB
Primary Accession	P11802
Other Accession	NM_000075.2
Reactivity	Human
Host	Mouse
Clonality	Monoclonal
Isotype	IgG1
Immunogen	Purified recombinant human CDK4 protein expressed in E.coli.
Purification	Affinity purified
Calculated MW	34kDa KDa

Anti-CDK4 Antibody - Additional Information

Gene ID 1019

Other Names

Cdk4;cdk4;CDK4protein;CDK4_HUMAN;Cell divisionkinase4;Celldivisionproteinkinase4;CMM3;CMM3;Crk3;Cyclindependentkinase4;Cyclin-dependentkinase4;Melanomacutaneousmalignant3;MGC14458;p34cdk4;PSKJ3;PSK-J3.

Dilution

WB~~1:500

Format


Purified mouse monoclonal antibody in PBS(pH 7.4) containing with 0.09% (W/V) sodium azide and 50% glycerol.

Storage

Store at -20 °C.Stable for 12 months from date of receipt

Anti-CDK4 Antibody - Protein Information

Name CDK4



Western blot detection of CDK4 in HeLa and K562 cell lysates using CDK4 mouse mAb (1:500 diluted).Predicted band size:34KDa.Observed band size:34KDa.

Anti-CDK4 Antibody - Background

Ser/Thr-kinase component of cyclin D-CDK4 (DC) complexes that phosphorylate and inhibit members of the retinoblastoma (RB) protein family including RB1 and regulate the cell-cycle during G(1)/S transition. Phosphorylation of RB1 allows dissociation of the

Function

Ser/Thr-kinase component of cyclin D-CDK4 (DC) complexes that phosphorylate and inhibit members of the retinoblastoma (RB) protein family including RB1 and regulate the cell-cycle during G(1)/S transition. Phosphorylation of RB1 allows dissociation of the transcription factor E2F from the RB/E2F complexes and the subsequent transcription of E2F target genes which are responsible for the progression through the G(1) phase. Hypophosphorylates RB1 in early G(1) phase. Cyclin D-CDK4 complexes are major integrators of various mitogenic and antimitogenic signals. Also phosphorylates SMAD3 in a cell-cycle-dependent manner and represses its transcriptional activity. Component of the ternary complex, cyclin D/CDK4/CDKN1B, required for nuclear translocation and activity of the cyclin D-CDK4 complex.

Cellular Location

Cytoplasm. Nucleus. Nucleus membrane. Note=Cytoplasmic when non-complexed. Forms a cyclin D-CDK4 complex in the cytoplasm as cells progress through G(1) phase. The complex accumulates on the nuclear membrane and enters the nucleus on transition from G(1) to S phase. Also present in nucleoli and heterochromatin lumps. Colocalizes with RB1 after release into the nucleus.

Anti-CDK4 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)