

REG3G Antibody (Center)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP5606c

Specification

REG3G Antibody (Center) - Product Information

Application	WB, IHC-P, FC,E
Primary Accession	Q6UW15
Other Accession	NP_001008388.1
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Antigen Region	89-116

REG3G Antibody (Center) - Additional Information

Gene ID 130120

Other Names

Regenerating islet-derived protein 3-gamma, REG-3-gamma, Pancreatitis-associated protein 1B, PAP-1B, Pancreatitis-associated protein IB, PAP IB, Regenerating islet-derived protein III-gamma, REG III, Reg III-gamma, Regenerating islet-derived protein 3-gamma 165 kDa form, Regenerating islet-derived protein 3-gamma 15 kDa form, REG3G, PAP1B

Target/Specificity

This REG3G antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 89-116 amino acids of human REG3G.

Dilution


WB~~1:1000
IHC-P~~1:25
FC~~1:10~50

Format


Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

Storage

Maintain refrigerated at 2-8°C for up to 2



All lanes : Anti-REG3G Antibody (Center) at 1:1000 dilution Lane 1: mouse small intestine tissue lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/15000 dilution. Observed band size : 20kDa Blocking/Dilution buffer: 5% NFDM/TBST.



Western blot analysis of lysate from SH-SY5Y cell line, using REG3G Antibody (Center)(Cat. #AP5606c). AP5606c was diluted at 1:1000. A goat anti-rabbit IgG H&L(HRP) at 1:5000

weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

REG3G Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

REG3G Antibody (Center) - Protein Information

Name REG3G

Synonyms PAP1B

Function

Bactericidal C-type lectin which acts exclusively against Gram-positive bacteria and mediates bacterial killing by binding to surface-exposed carbohydrate moieties of peptidoglycan. Restricts bacterial colonization of the intestinal epithelial surface and consequently limits activation of adaptive immune responses by the microbiota. The uncleaved form has bacteriostatic activity, whereas the cleaved form has bactericidal activity against *L.monocytogenes* and methicillin-resistant *S.aureus*. Regulates keratinocyte proliferation and differentiation after skin injury.

Cellular Location

Secreted. Cytoplasm.

Tissue Location


Predominantly expressed in pancreas, where it may be restricted to exocrine pancreas. Moderate expression levels in testis and weak in heart, kidney and placenta

REG3G Antibody (Center) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)

dilution was used as the secondary antibody. Lysate at 35ug.




REG3G Antibody (Center) (Cat.# AP5606c) western blot analysis in SH-SY5Y cell line lysates (35ug/lane). This demonstrates the REG3G antibody detected the REG3G protein (arrow).




All lanes : Anti-REG3G Antibody (Center) at 1:100 dilution
Lane 1: human kidney lysates
Lane 2: human pancreas lysates
Lysates/proteins at 20 µg per lane.
Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution
Predicted band size : 19 kDa
Blocking/Dilution buffer: 5% NFDM/TBST.


- [Flow Cytometry](#)
- [Cell Culture](#)




Anti-REG3G Antibody (Center) at 1:1000 dilution + recombinant protein lysates
Lysates/proteins at 20 µg per lane.
Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution
Predicted band size : 19 kDa
Blocking/Dilution buffer: 5% NFDM/TBST.




Anti-REG3G Antibody (Center) at 1:400 dilution + human kidney lysates
Lysates/proteins at 20 µg per lane.
Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution
Predicted band size : 19 kDa
Blocking/Dilution buffer: 5% NFDM/TBST.




All lanes : Anti-REG3G Antibody (Center) at 1:2000 dilution
Lane 1: human kidney lysates
Lane 2: human lung lysates
Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 19 kDa
Blocking/Dilution buffer: 5% NFDM/TBST.




Anti-REG3G Antibody (Center) at 1:2000 dilution + human pancreas lysate
Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 19 kDa
Blocking/Dilution buffer: 5% NFDM/TBST.




Anti-REG3G Antibody (Center) at 1:1000 dilution + human kidney lysates
Lysates/proteins at 20 µg per lane.
Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution.
Predicted band size : 19 kDa
Blocking/Dilution buffer: 5% NFDM/TBST.



REG3G Antibody (Center) (Cat. #AP5606c) immunohistochemistry analysis in formalin fixed and paraffin embedded human pancreas tissue followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of the REG3G Antibody (Center) for immunohistochemistry. Clinical relevance has not been evaluated.




AP5606C staining REG3G in human small intestine sections by Immunohistochemistry (IHC-P - paraformaldehyde-fixed, paraffin-embedded sections). Tissue was fixed with formaldehyde and blocked with 3% BSA for 0.5 hour at room temperature; antigen retrieval was by heat mediation with a citrate buffer (pH6). Samples were incubated with primary antibody (1/25) for 1 hour at 37°C. A undiluted biotinylated goat polyvalent antibody was used as the secondary antibody.



AP5606C staining REG3G in human small intestine sections by Immunohistochemistry (IHC-P - paraformaldehyde-fixed, paraffin-embedded sections). Tissue was fixed with formaldehyde and blocked with 3% BSA for 0.5 hour at room temperature; antigen retrieval was by heat mediation with a citrate buffer (pH6). Samples were incubated with primary antibody (1/25) for 1

hours at 37°C. A undiluted biotinylated goat polyvalent antibody was used as the secondary antibody.

Abgent antibody neutralizes Reg3 γ binding to *S. aureus*



Reg3g binds to Staphylococcus aureus and the antibody did block some of this binding (Kindly offered by Dr. Choi).

REG3G Antibody (Center) - Background

REG3G might be a stress protein involved in the control of bacterial proliferation (By similarity).

REG3G Antibody (Center) - References

Taylor-Fishwick, D.A., et al. J. Histochem. Cytochem. 56(2):183-191(2008)
Hamblet, N.S., et al. Pancreas 36(1):1-9(2008)
Hillier, L.W., et al. Nature 434(7034):724-731(2005)
Laurine, E., et al. Biochim. Biophys. Acta 1727(3):177-187(2005)

REG3G Antibody (Center) - Citations

- [The colonic macrophage transcription factor RBP-J orchestrates intestinal immunity against bacterial pathogens.](#)
- [Commensal bacteria direct selective cargo sorting to promote symbiosis.](#)
- [Reg3g Promotes Pancreatic Carcinogenesis in a Murine Model of Chronic Pancreatitis.](#)
- [Autophagy in pancreatic acinar cells in caerulein-treated mice: immunolocalization of related proteins and their potential as markers of pancreatitis.](#)
- [Innate Stat3-mediated induction of the antimicrobial protein Reg3 \$\gamma\$ is required for host defense against MRSA pneumonia.](#)