

**MRPL47 Polyclonal Antibody**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP56837****Specification****MRPL47 Polyclonal Antibody - Product Information**

Application	<b>IHC-P</b>
Primary Accession	<a href="#">Q9HD33</a>
Reactivity	<b>Rat, Pig, Dog, Cow</b>
Host	<b>Rabbit</b>
Clonality	<b>Polyclonal</b>
Calculated MW	<b>29450</b>

**MRPL47 Polyclonal Antibody - Additional Information****Gene ID** 57129**Other Names**

39S ribosomal protein L47, mitochondrial, L47mt, MRP-L47, Mitochondrial large ribosomal subunit protein uL29m, Nasopharyngeal carcinoma metastasis-related protein 1, MRPL47, NCM1

**Format**


0.01M TBS(pH7.4) with 1% BSA, 0.09% (W/V) sodium azide and 50% Glyce

**Storage**

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

**MRPL47 Polyclonal Antibody - Protein Information****Name** MRPL47**Synonyms** NCM1**Cellular Location**

Mitochondrion



Paraformaldehyde-fixed, paraffin embedded (Mouse brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (MRPL47) Polyclonal Antibody, Unconjugated (bs-17800R) at 1:500 overnight at 4°C, followed by a conjugated secondary (sp-0023) for 20 minutes and DAB staining.

## **MRPL47 Polyclonal Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)