

NAGA Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP57346**Specification****NAGA Polyclonal Antibody - Product Information**

Application	IHC-P
Primary Accession	P17050
Reactivity	Rat, Pig, Dog, Cow
Host	Rabbit
Clonality	Polyclonal
Calculated MW	46565

NAGA Polyclonal Antibody - Additional Information**Gene ID 4668****Other Names**

Alpha-N-acetylgalactosaminidase, 3.2.1.49, Alpha-galactosidase B, NAGA (http://www.genenames.org/cgi-bin/gene_symbol_report?hgnc_id=7631)
target="_blank">HGNC:7631)

Format

0.01M TBS(pH7.4) with 1% BSA, 0.09% (W/V) sodium azide and 50% Glycerol

Storage


Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

NAGA Polyclonal Antibody - Protein Information**Name** NAGA ([HGNC:7631](#))**Function**

Removes terminal alpha-N-acetylgalactosamine residues from glycolipids and glycopeptides. Required for the breakdown of glycolipids.

Cellular Location

Lysosome.



Paraformaldehyde-fixed, paraffin embedded (Rat liver); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (NAGA) Polyclonal Antibody, Unconjugated (bs-19001R) at 1:500 overnight at 4°C, followed by a conjugated secondary (sp-0023) for 20 minutes and DAB staining.

NAGA Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)