

**CYP4A22 Polyclonal Antibody**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP58268**

**Specification**

**CYP4A22 Polyclonal Antibody - Product Information**

Application	<b>WB</b>
Primary Accession	<a href="#">Q5TCH4</a>
Reactivity	<b>Rat</b>
Host	<b>Rabbit</b>
Clonality	<b>Polyclonal</b>
Calculated MW	<b>59246</b>

**CYP4A22 Polyclonal Antibody - Additional Information**

**Gene ID** 284541

**Other Names**

Cytochrome P450 4A22, CYP4A22, Fatty acid omega-hydroxylase, Lauric acid omega-hydroxylase, Long-chain fatty acid omega-monooxygenase, 1.14.14.80, CYP4A22

**Format**

0.01M TBS(pH7.4) with 1% BSA, 0.09% (W/V) sodium azide and 50% Glyce

**Storage**


Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

**CYP4A22 Polyclonal Antibody - Protein Information**

**Name** CYP4A22

**Function**

Catalyzes the omega- and (omega-1)-hydroxylation of various fatty acids such as laurate and palmitate. Shows no activity towards arachidonic acid and prostaglandin A1. Lacks functional activity in the kidney and does not contribute to renal 20-hydroxyecosatetraenoic acid



**Sample:**  
Liver (Mouse) Lysate at 40 ug  
Spleen (Mouse) Lysate at 40 ug  
Primary: Anti-CYP4A22 (bs-5055R) at 1/1000 dilution  
Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution  
Predicted band size: 58 kD  
Observed band size: 58 kD

(20-HETE) biosynthesis.

**Cellular Location**

Endoplasmic reticulum membrane;  
Peripheral membrane protein. Microsome  
membrane; Peripheral membrane protein

**CYP4A22 Polyclonal Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)