

FAM13C1 Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP58919

Specification

FAM13C1 Polyclonal Antibody - Product Information

Application	IHC-P, WB
Primary Accession	Q8NE31
Reactivity	Rat, Pig
Host	Rabbit
Clonality	Polyclonal
Calculated MW	65727

FAM13C1 Polyclonal Antibody - Additional Information

Gene ID 220965

Other Names

Protein FAM13C, FAM13C, FAM13C1

Format

0.01M TBS(pH7.4) with 1% BSA, 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

FAM13C1 Polyclonal Antibody - Protein Information


Name FAM13C

Synonyms FAM13C1


FAM13C1 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)




Tissue/cell: Rat brain tissue; 4% Paraformaldehyde-fixed and paraffin-embedded;
Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at 37°C for 20 min;
Incubation: Anti-FAM13C1 Polyclonal Antibody, Unconjugated(bs-8215R) 1:500, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining




Paraformaldehyde-fixed, paraffin embedded (rat lung); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3%

- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (FAM13C1) Polyclonal Antibody, Unconjugated (bs-8215R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.




Paraformaldehyde-fixed, paraffin embedded (mouse liver); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (FAM13C1) Polyclonal Antibody, Unconjugated (bs-8215R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Sample: MCF-7 Cell (Human) Lysate at 40 ug
Primary: Anti-FAM13C1 (bs-8215R) at 1/300 dilution
Secondary: IRDye800CW Goat Anti-Rabbit

IgG at 1/20000 dilution
Predicted band size: 66 kD
Observed band size: 66 kD



Sample:
Hela Cell (Human) Lysate at 30 ug
Primary: Anti- FAM13C1 (bs-8215R) at 1/300 dilution
Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution
Predicted band size: 66 kD
Observed band size: 63 kD