

TIM 3 Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP59082

Specification

TIM 3 Polyclonal Antibody - Product Information

Application	WB, IHC-P
Reactivity	Rat
Host	Rabbit
Clonality	Polyclonal

TIM 3 Polyclonal Antibody - Additional Information

Format

0.01M TBS(pH7.4) with 1% BSA, 0.09% (W/V) sodium azide and 50% Glyce

Storage


Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

TIM 3 Polyclonal Antibody - Protein Information


TIM 3 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)




Sample:
Thymus (Mouse) Lysate at 40 ug
Raw264.7(Mouse) Cell Lysate at 30 ug
Primary: Anti-TIM 3 (bs-8766R) at 1/1000 dilution
Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution
Predicted band size: 29 kD
Observed band size: 49 kD




Sample:

Liver(Rat) Lysate at 40 ug
Primary: Anti-TIM 3 (bs-8766R) at 1/1000 dilution
Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution
Predicted band size: 29 kD
Observed band size: 49 kD



Sample:
Spleen (Mouse) Lysate at 40 ug
Primary: Anti-TIM 3 (bs-8766R) at 1/1000 dilution
Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution
Predicted band size: 29 kD
Observed band size: 49 kD



Tissue/cell: rat liver tissue; 4% Paraformaldehyde-fixed and paraffin-embedded;
Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal

goat serum,C-0005) at 37°C for 20 min;
Incubation: Anti-TIM-3 Polyclonal Antibody,
Unconjugated(bs-8766R) 1:200, overnight at
4°C, followed by conjugation to the
secondary antibody(SP-0023) and
DAB(C-0010) staining