

MAP7D1 Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP59217

Specification

MAP7D1 Polyclonal Antibody - Product Information

Application	WB, IHC-P
Primary Accession	Q3KQU3
Reactivity	Rat, Pig, Dog, Cow
Host	Rabbit
Clonality	Polyclonal
Calculated MW	92820

MAP7D1 Polyclonal Antibody - Additional Information

Gene ID 55700

Other Names

MAP7 domain-containing protein 1, Arginine/proline-rich coiled-coil domain-containing protein 1, Proline/arginine-rich coiled-coil domain-containing protein 1, MAP7D1, KIAA1187, PARCC1, RPRC1

Format

0.01M TBS(pH7.4) with 1% BSA, 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.


MAP7D1 Polyclonal Antibody - Protein Information

Name MAP7D1

Synonyms KIAA1187, PARCC1, RPRC1

Cellular Location

Cytoplasm, cytoskeleton, spindle




Sample:

Heart (Mouse) Lysate at 40 ug
Primary: Anti- MAP7D1 (bs-9314R) at 1/1000 dilution

Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution

Predicted band size: 93 kD

Observed band size: 93 kD



Paraformaldehyde-fixed, paraffin embedded (mouse brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (MAP7D1)

MAP7D1 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Polyclonal Antibody, Unconjugated (bs-9314R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.