

LRP1 Antibody (C-term)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP6153a

Specification

LRP1 Antibody (C-term) - Product Information

Application	IHC-P,E
Primary Accession	Q07954
Other Accession	Q91ZX7 , G3V928
Reactivity	Human
Predicted	Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit Ig
Antigen Region	4515-4544

LRP1 Antibody (C-term) - Additional Information

Gene ID 4035

Other Names

Prolow-density lipoprotein receptor-related protein 1, LRP-1, Alpha-2-macroglobulin receptor, A2MR, Apolipoprotein E receptor, APOER, CD91, Low-density lipoprotein receptor-related protein 1 85 kDa subunit, LRP-85, Low-density lipoprotein receptor-related protein 1 515 kDa subunit, LRP-515, Low-density lipoprotein receptor-related protein 1 intracellular domain, LRPICD, LRP1, A2MR, APR

Target/Specificity

This LRP1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 4515~4544 amino acids from the C-terminal region of human LRP1.

Dilution


IHC-P~~1:50~100

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

Storage

Maintain refrigerated at 2-8°C for up to 2



Formalin-fixed and paraffin-embedded human cancer tissue reacted with the primary antibody, which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated. BC = breast carcinoma; HC = hepatocarcinoma.

LRP1 Antibody (C-term) - Background

Low density lipoprotein (LDL) receptor-related protein (LRP), a member of the LDL receptor family, binds multiple classes of ligands and has been implicated in a broad range of normal and disease processes involving lipid metabolism, protease clearance, and cell migration (1). Structurally, members of the LDLR family share homology within their extracellular domains, which are highlighted by the presence of clusters of ligand-binding repeats. LRP is a large endocytic receptor that participates in several biological pathways and plays prominent roles in lipoprotein metabolism and in the catabolism of proteinases involved in coagulation and fibrinolysis. LRP also mediates the cellular entry of certain viruses and toxins and facilitates the activation of various lysosomal

weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

LRP1 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

LRP1 Antibody (C-term) - Protein Information

Name LRP1 ([HGNC:6692](#))

Synonyms A2MR, APR

Function

Endocytic receptor involved in endocytosis and in phagocytosis of apoptotic cells (PubMed:11907044, PubMed:12713657). Required for early embryonic development (By similarity). Involved in cellular lipid homeostasis. Involved in the plasma clearance of chylomicron remnants and activated LRPAP1 (alpha 2-macroglobulin), as well as the local metabolism of complexes between plasminogen activators and their endogenous inhibitors. Acts as an LRPAP1 alpha-2- macroglobulin receptor (PubMed:26142438, PubMed:1702392). Acts as TAU/MAPT receptor and controls the endocytosis of TAU/MAPT as well as its subsequent spread (PubMed:32296178). May modulate cellular events, such as APP metabolism, kinase-dependent intracellular signaling, neuronal calcium signaling as well as neurotransmission (PubMed:12888553).

Cellular Location

[Low-density lipoprotein receptor-related protein 1 85 kDa subunit]: Cell membrane; Single-pass type I membrane protein Membrane, coated pit [Low-density

enzymes (2). All LRPs are expressed in the central nervous system and, for most receptors, animal models have shown that they are indispensable for successful neurodevelopment. The mechanisms by which they regulate the formation of the nervous system are varied and include the transduction of extracellular signals and the modulation of intracellular signal propagation, as well as cargo transport, the function most commonly attributed to this gene family (3).

LRP1 Antibody (C-term) - References

Kinoshita, A., et al., J. Biol. Chem. 278(42):41182-41188 (2003).
Zhu, Y., et al., J. Biol. Chem. 278(38):36257-36263 (2003).
Lutters, B.C., et al., J. Biol. Chem. 278(36):33831-33838 (2003).
Stebbing, J., et al., Blood 102(5):1806-1814 (2003).
Takayama, Y., et al., J. Biol. Chem. 278(24):22112-22118 (2003).

lipoprotein receptor-related protein 1
intracellular domain]: Cytoplasm Nucleus.

Note=After cleavage, the intracellular
domain (LRPICD) is detected both in the
cytoplasm and in the nucleus.

Tissue Location

Most abundant in liver, brain and lung.

LRP1 Antibody (C-term) - Protocols

Provided below are standard protocols that you
may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)