

NRG3 Antibody (Center)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP6224a

Specification

NRG3 Antibody (Center) - Product Information

Application	WB, IHC-P,E
Primary Accession	P56975
Other Accession	O35181
Reactivity	Human
Predicted	Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit Ig
Antigen Region	305-336

NRG3 Antibody (Center) - Additional Information

Gene ID 10718

Other Names

Pro-neuregulin-3, membrane-bound isoform, Pro-NRG3, Neuregulin-3, NRG-3, NRG3

Target/Specificity

This NRG3 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 305-336 amino acids from the Central region of human NRG3.

Dilution

WB~~1:1000

IHC-P~~1:50~100

Format


Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

Storage


Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

NRG3 Antibody (Center) is for research use



NRG3 Antibody (Center)(Cat. #AP6224a) western blot analysis in human blood plasma tissue lysates (35ug/lane).This demonstrates the NRG3 antibody detected the NRG3 protein (arrow).



Formalin-fixed and paraffin-embedded human cancer tissue reacted with the primary antibody, which was peroxidase-conjugated to the secondary antibody, followed by AEC staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated. BC = breast carcinoma;

only and not for use in diagnostic or therapeutic procedures.

HC = hepatocarcinoma.

NRG3 Antibody (Center) - Protein Information

Name NRG3

Function

Direct ligand for the ERBB4 tyrosine kinase receptor. Binding results in ligand-stimulated tyrosine phosphorylation and activation of the receptor. Does not bind to the EGF receptor, ERBB2 or ERBB3 receptors. May be a survival factor for oligodendrocytes.

Cellular Location

[Pro-neuregulin-3, membrane-bound isoform]: Cell membrane; Single-pass type I membrane protein. Note=Does not seem to be active. [Isoform 3]: Cell membrane; Single-pass type I membrane protein. Note=Isoform 3 is also proteolytically released as a soluble form

Tissue Location

Highly expressed in most regions of the brain with the exception of corpus callosum. Expressed at lower level in testis Not detected in heart, placenta, lung, liver, skeletal muscle, kidney, pancreas, spleen, thymus, prostate, ovary, small intestine, colon and peripheral blood leukocytes

NRG3 Antibody (Center) - Background

NRG3, which belongs to the neuregulin family, is a direct ligand for the ERBB4 tyrosine kinase receptor. Binding results in ligand-stimulated tyrosine phosphorylation and activation of the receptor. NRG3 does not bind to the EGF receptor, ERBB2 or ERBB3 receptors. The protein exists as an type I membrane protein and as a proteolytically released soluble growth factor form. The membrane-bound form does not appear to be active. NRG3 is highly expressed in most regions of the brain with the exception of corpus callosum, and is expressed at lower level in testis. It is not detected in heart, placenta, lung, liver, skeletal muscle, kidney, pancreas, spleen, thymus, prostate, ovary, small intestine, colon and peripheral blood leukocytes. The NRG3 cytoplasmic domain may be involved in the regulation of trafficking and proteolytic processing. Regulation of the proteolytic processing may involve initial intracellular domain dimerization.

NRG3 Antibody (Center) - References

Zhang, D., et al., Proc. Natl. Acad. Sci. U.S.A. 94(18):9562-9567 (1997).

NRG3 Antibody (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

NRG3 Antibody (Center) - Citations

- [Levels of neuregulin 1 and 3 proteins in Brodmann's area 46 from subjects with schizophrenia and bipolar disorder.](#)
- [Identification of the scaramanga gene implicates Neuregulin3 in mammary gland specification.](#)

