

Presenilin 1 (PSEN1) Antibody (C-term)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP6231A

Specification

Presenilin 1 (PSEN1) Antibody (C-term) - Product Information

Application	IF, WB, IHC-P, FC, E
Primary Accession	P49768
Other Accession	P97887 , P49769 , Q8HXW5
Reactivity	Human, Mouse
Predicted	Monkey, Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit Ig
Antigen Region	330-359

Presenilin 1 (PSEN1) Antibody (C-term) - Additional Information

Gene ID 5663

Other Names

Presenilin-1, PS-1, 3423-, Protein S182, Presenilin-1 NTF subunit, Presenilin-1 CTF subunit, Presenilin-1 CTF12, PS1-CTF12, PSEN1, AD3, PS1, PSNL1

Target/Specificity

This Presenilin 1 (PSEN1) antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 330-359 amino acids from the C-terminal region of human Presenilin 1 (PSEN1).


Dilution

IF~~1:10~50
WB~~1:1000
IHC-P~~1:50~100
FC~~1:10~50


Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

Storage



Confocal immunofluorescent analysis of Presenilin 1 (PSEN1) Antibody (C-term)(Cat#AP6231a) with MDA-MB435 cell followed by Alexa Fluor 488-conjugated goat anti-rabbit IgG (green). DAPI was used to stain the cell nuclear (blue).



All lanes : Anti-PSEN1 Antibody (C-term) at 1:1000 dilution
Lane 1: HeLa whole cell lysate
Lane 2: HL-60 whole cell lysate
Lysates/proteins at 20 µg per lane.
Secondary Goat Anti-Rabbit IgG, (H+L),

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Presenilin 1 (PSEN1) Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Presenilin 1 (PSEN1) Antibody (C-term) - Protein Information

Name PSEN1

Synonyms AD3, PS1, PSNL1

Function

Catalytic subunit of the gamma-secretase complex, an endoprotease complex that catalyzes the intramembrane cleavage of integral membrane proteins such as Notch receptors and APP (amyloid- beta precursor protein) (PubMed:15274632,

PubMed:10545183,

PubMed:10593990,

PubMed:10206644,

PubMed:10899933,

PubMed:10811883,

PubMed:12679784,

PubMed:12740439,


PubMed:25043039,

PubMed:26280335,


PubMed:30598546,

target="_blank">30598546,

Peroxidase conjugated at 1/10000 dilution.
Predicted band size : 53 kDa
Blocking/Dilution buffer: 5% NFDM/TBST.



Formalin-fixed and paraffin-embedded human cancer tissue reacted with the primary antibody, which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated. BC = breast carcinoma; HC = hepatocarcinoma.



Presenilin 1 (PSEN1) Antibody (C-term) (Cat. #AP6231a) flow cytometric analysis of Hela cells (right histogram) compared to a negative control cell (left histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

Presenilin 1 (PSEN1) Antibody (C-term) - Background

Alzheimer's disease (AD) patients with an inherited form of the disease carry mutations

PubMed:30630874, PubMed:28269784, PubMed:20460383). Requires the presence of the other members of the gamma-secretase complex for protease activity (PubMed:15274632, PubMed:25043039, PubMed:26280335, PubMed:30598546, PubMed:30630874). Plays a role in Notch and Wnt signaling cascades and regulation of downstream processes via its role in processing key regulatory proteins, and by regulating cytosolic CTNNB1 levels (PubMed:9738936, PubMed:10593990, PubMed:10899933, PubMed:10811883). Stimulates cell-cell adhesion via its interaction with CDH1; this stabilizes the complexes between CDH1 (E-cadherin) and its interaction partners CTNNB1 (beta-catenin), CTNND1 and JUP (gamma-catenin) (PubMed:11953314). Under conditions of apoptosis or calcium influx, cleaves CDH1 (PubMed:11953314). This promotes the disassembly of the complexes between CDH1 and CTNND1, JUP and CTNNB1, increases the pool of cytoplasmic

in the presenilin proteins (PSEN1; PSEN2) or the amyloid precursor protein (APP). These disease-linked mutations result in increased production of the longer form of amyloid-beta (main component of amyloid deposits found in AD brains). Presenilins are postulated to regulate APP processing through their effects on gamma-secretase, an enzyme that cleaves APP. Also, it is thought that the presenilins are involved in the cleavage of the Notch receptor, such that they either directly regulate gamma-secretase activity or themselves are protease enzymes.

Presenilin 1 (PSEN1) Antibody (C-term) - References

Marambaud, P., et al., Cell 114(5):635-645 (2003). Kim, S.H., et al., J. Biol. Chem. 278(36):33992-34002 (2003). Miklossy, J., et al., Neurobiol. Aging 24(5):655-662 (2003). Cai, D., et al., J. Biol. Chem. 278(5):3446-3454 (2003). Godin, C., et al., Neuroreport 14(12):1613-1616 (2003).

CTNNB1, and thereby negatively regulates Wnt signaling (PubMed:9738936, PubMed:11953314). Required for normal embryonic brain and skeleton development, and for normal angiogenesis (By similarity). Mediates the proteolytic cleavage of EphB2/CTF1 into EphB2/CTF2 (PubMed:17428795, PubMed:28269784). The holoprotein functions as a calcium-leak channel that allows the passive movement of calcium from endoplasmic reticulum to cytosol and is therefore involved in calcium homeostasis (PubMed:25394380, PubMed:16959576). Involved in the regulation of neurite outgrowth (PubMed:15004326, PubMed:20460383). Is a regulator of presynaptic facilitation, spike transmission and synaptic vesicles replenishment in a process that depends on gamma-secretase activity. It acts through the control of SYT7 presynaptic expression (By similarity).

Cellular Location

Endoplasmic reticulum. Endoplasmic reticulum membrane; Multi-pass membrane protein. Golgi apparatus membrane; Multi-pass membrane protein. Cytoplasmic granule. Cell membrane; Multi-pass membrane protein. Cell projection, growth cone. Early endosome. Early endosome membrane; Multi-pass membrane protein. Cell projection, neuron projection. Cell projection, axon
{ECO:0000250|UniProtKB:Q4JIM4}. Cell junction, synapse
{ECO:0000250|UniProtKB:Q4JIM4}.
Note=Translocates with bound NOTCH1 from the endoplasmic reticulum and/or

Golgi to the cell surface (PubMed:10593990). Colocalizes with CDH1/2 at sites of cell-cell contact. Colocalizes with CTNNB1 in the endoplasmic reticulum and the proximity of the plasma membrane (PubMed:9738936). Also present in azurophil granules of neutrophils (PubMed:11987239). Colocalizes with UBQLN1 in the cell membrane and in cytoplasmic juxtanuclear structures called aggresomes (PubMed:21143716).

Tissue Location

Detected in azurophile granules in neutrophils and in platelet cytoplasmic granules (at protein level) (PubMed:11987239) Expressed in a wide range of tissues including various regions of the brain, liver, spleen and lymph nodes (PubMed:7596406, PubMed:8641442, PubMed:8574969).

Presenilin 1 (PSEN1) Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Presenilin 1 (PSEN1) Antibody (C-term) - Citations

- [Nicotine decreases beta-amyloid through regulating BACE1 transcription in SH-EP1- \$\Delta\$ 4 \$\Delta\$ 2 nAChR-APP695 cells.](#)