

INA (alpha internexin) Antibody (Center)

Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP6284c

Specification

INA (alpha internexin) Antibody (Center) - Product Information

Application WB, IHC-P,E Primary Accession 016352

Other Accession P23565, P46660,

Q08DH7

Reactivity Human, Mouse,

Rat

Predicted Bovine
Host Rabbit
Clonality Polyclonal
Isotype Rabbit IgG
Antigen Region 290-319

INA (alpha internexin) Antibody (Center) - Additional Information

Gene ID 9118

Other Names

Alpha-internexin, Alpha-Inx, 66 kDa neurofilament protein, NF-66, Neurofilament-66, Neurofilament 5, INA, NFF5

Target/Specificity

This INA (alpha internexin) antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 290-319 amino acids from the Central region of human INA (alpha internexin).

Dilution


WB~~1:2000 IHC-P~~1:10~50

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.


Storage


Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C



All lanes: Anti-INA Antibody (Center) at 1:1000-2000 dilution Lane 1: Human brain tissue lysate Lane 2: U-87 MG whole cell lysate Lane 3: SH-SY5Y whole cell lysate Lane 4: Mouse liver tissue lysate Lane 5: Mouse brain tissue lysate Lane 6: Rat brain tissue lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size: 55 kDa

Blocking/Dilution buffer: 5% NFDM/TBST.





Tel: 858.875.1900 Fax: 858.622.0609



in small aliquots to prevent freeze-thaw cycles.

Precautions

INA (alpha internexin) Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

INA (alpha internexin) Antibody (Center) - Protein Information

Name INA

Synonyms NEF5

Function

Class-IV neuronal intermediate filament that is able to self- assemble. It is involved in the morphogenesis of neurons. It may form an independent structural network without the involvement of other neurofilaments or it may cooperate with NEFL to form the filamentous backbone to which NEFM and NEFH attach to form the cross-bridges. May also cooperate with the neuronal intermediate filament protein PRPH to form filamentous networks (By similarity).

Tissue Location

Found predominantly in adult CNS.

INA (alpha internexin) Antibody (Center) -**Protocols**

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- Immunoprecipitation
- Flow Cvtometv
- Cell Culture

All lanes: Anti-INA Antibody (Center) at 1:2000 dilution Lane 1: Human brain tissue lysate Lane 2: Mouse brain tissue lysate Lane 3: Rat brain tissue lysate Lane 4: SH-SY5Y whole cell lysate Lane 5: Human liver tissue lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size: 55 kDa Blocking/Dilution buffer: 5% NFDM/TBST.




Formalin-fixed and paraffin-embedded human brain tissue reacted with INA antibody (Center) (Cat.#AP6284c), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.

INA (alpha internexin) Antibody (Center) -**Background**

INA is a class-IV neuronal intermediate filament that is able to self-assemble. It is involved in the morphogenesis of neurons. It may form an independent structural network without the involvement of other neurofilaments or it may cooperate with NF-L to form the filamentous backbone to which NF-M and NF-H attach to form the cross-bridges.

INA (alpha internexin) Antibody (Center) -References

Armstrong, R.A., Eur. J. Neurol. 13 (5), 528-532 (2006)Suzuki, T., Eur. J. Neurosci. 21 (2), 339-350 Cairns, N.J., Am. J. Pathol. 164 (6), 2153-2161 (2004)





 INA (alpha internexin) Antibody (Center) - Citations
 EGFR Amplification and IDH Mutations in Glioblastoma Patients of the Northeast of Morocco.