

**NUR77 (NR4A1) Antibody (S351)**  
**Affinity Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP6289c**

**Specification**

**NUR77 (NR4A1) Antibody (S351) - Product Information**

Application	WB,E
Primary Accession	<a href="#">P22736</a>
Other Accession	<a href="#">P22829</a> , <a href="#">P12813</a>
Reactivity	Human, Mouse, Rat
Predicted Host	Mouse, Rat
Clonality	Rabbit
Isotype	Polyclonal
Antigen Region	Rabbit Ig 329-358

**NUR77 (NR4A1) Antibody (S351) - Additional Information**

**Gene ID** 3164

**Other Names**

Nuclear receptor subfamily 4 group A member 1, Early response protein NAK1, Nuclear hormone receptor NUR/77, Nur77, Orphan nuclear receptor HMR, Orphan nuclear receptor TR3, ST-59, Testicular receptor 3, NR4A1, GFRP1, HMR, NAK1

**Target/Specificity**

This NUR77(NR4A1) antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 329-358 amino acids from human NUR77(NR4A1).

**Dilution**


WB~~1:500

**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C



All lanes : Anti-NUR77 (NR4A1) Antibody (S351) at 1:500 dilution Lane 1: Jurkat whole cell lysate Lane 2: Hela whole cell lysate Lane 3: Human liver tissue lysate Lane 4: MCF-7 whole cell lysate Lane 5: Mouse brain tissue lysate Lane 6: NIH/3T3 whole cell lysate Lane 7: Rat brain tissue lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 64 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

**NUR77 (NR4A1) Antibody (S351) - Background**

NR4A1 is a member of the steroid-thyroid hormone-retinoid receptor superfamily. Expression is induced by phytohemagglutinin in human lymphocytes and by serum stimulation of arrested fibroblasts. This protein acts as a nuclear transcription factor. Translocation of the protein from the nucleus to mitochondria induces apoptosis.

in small aliquots to prevent freeze-thaw cycles.

**Precautions**

NUR77 (NR4A1) Antibody (S351) is for research use only and not for use in diagnostic or therapeutic procedures.

**NUR77 (NR4A1) Antibody (S351) - Protein Information**

**Name** NR4A1

**Synonyms** GFRP1, HMR, NAK1

**Function**

Orphan nuclear receptor. May act concomitantly with NURR1 in regulating the expression of delayed-early genes during liver regeneration. Binds the NGFI-B response element (NBRE) 5'-AAAAGGTCA-3' (By similarity). May inhibit NF-kappa-B transactivation of IL2. Participates in energy homeostasis by sequestering the kinase STK11 in the nucleus, thereby attenuating cytoplasmic AMPK activation. Plays a role in the vascular response to injury (By similarity).

**Cellular Location**

Nucleus. Cytoplasm. Mitochondrion. Note=Nuclear export to the cytoplasm is XPO1-mediated and positively regulated by IFI27 (PubMed:22427340) Translocation to the mitochondrion upon interaction with RXRA and upon the presence of 9-cis retinoic acid (PubMed:17761950)

**Tissue Location**

Fetal muscle and adult liver, brain and thyroid.

**NUR77 (NR4A1) Antibody (S351) - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)

- [Flow Cytometry](#)
- [Cell Culture](#)

**NUR77 (NR4A1) Antibody (S351) - Citations**

- [Validation of an oligo-gene signature for the prognostic stratification of ductal carcinoma in situ \(DCIS\).](#)