

**VIP Antibody (C-term)**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP6554b**

**Specification**

**VIP Antibody (C-term) - Product Information**

Application	<b>WB, IHC-P, FC,E</b>
Primary Accession	<a href="#">P01282</a>
Reactivity	<b>Human</b>
Host	<b>Rabbit</b>
Clonality	<b>Polyclonal</b>
Isotype	<b>Rabbit Ig</b>
Calculated MW	<b>19169</b>
Antigen Region	<b>139-167</b>

**VIP Antibody (C-term) - Additional Information**

**Gene ID 7432**

**Other Names**

VIP peptides, Intestinal peptide PHV-42, Peptide histidine valine 42, Intestinal peptide PHM-27, Peptide histidine methioninamide 27, Vasoactive intestinal peptide, VIP, Vasoactive intestinal polypeptide, VIP

**Target/Specificity**

This VIP antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 139-167 amino acids from the C-terminal region of human VIP.

**Dilution**


WB~~1:1000  
IHC-P~~1:50~100  
FC~~1:10~50

**Format**


Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.



Western blot analysis of VIP antibody (C-term) (Cat. #AP6554b) in Jurkat cell line lysates (35ug/lane). VIP (arrow) was detected using the purified Pab.



Formalin-fixed and paraffin-embedded human prostate carcinoma with VIP Antibody (C-term), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.

### Precautions

VIP Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

### VIP Antibody (C-term) - Protein Information


#### Name VIP

#### Function

VIP causes vasodilation, lowers arterial blood pressure, stimulates myocardial contractility, increases glycogenolysis and relaxes the smooth muscle of trachea, stomach and gall bladder.

#### Cellular Location

Secreted.



Flow cytometric analysis of jurkat cells using VIP Antibody (C-term)(bottom histogram) compared to a negative control cell (top histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

### VIP Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### VIP Antibody (C-term) - Background

VIP belongs to the glucagon family. It stimulates myocardial contractility, causes vasodilation, increases glycogenolysis, lowers arterial blood pressure and relaxes the smooth muscle of trachea, stomach and gall bladder.

### VIP Antibody (C-term) - References

Lu,Y., Neurogastroenterol. Motil. 21 (7), 754-E47 (2009)  
Zhu,L., Int J Gynaecol Obstet 105 (3), 223-225 (2009)