

### SPP1 Antibody (C-term)

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP6555b

### **Specification**

#### SPP1 Antibody (C-term) - Product Information

Application
Primary Accession
Reactivity
Host
Clonality
Isotype
WB, IHC-P, FC,E
P10451
Human
Rabbit
Polyclonal
Rabbit Ig

SPP1 Antibody (C-term) - Additional Information

273-301

#### **Gene ID** 6696

Antigen Region

### **Other Names**

Osteopontin, Bone sialoprotein 1, Nephropontin, Secreted phosphoprotein 1, SPP-1, Urinary stone protein, Uropontin, SPP1, BNSP, OPN

# **Target/Specificity**

This SPP1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 273-301 amino acids from the C-terminal region of human SPP1.

### **Dilution**

WB~~1:2000 IHC-P~~1:50~100 FC~~1:10~50

#### **Format**


Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

### **Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.


# **Precautions**

SPP1 Antibody (C-term) is for research use



All lanes: Anti-SPP1 Antibody (C-term) at 1:2000 dilution Lane 1: Human plasma lysate Lane 2: Human placenta lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size: 35 kDa

Blocking/Dilution buffer: 5% NFDM/TBST.



Formalin-fixed and paraffin-embedded human lung carcinoma with SPP1 Antibody (C-term), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.



only and not for use in diagnostic or therapeutic procedures.

SPP1 Antibody (C-term) - Protein Information

Name SPP1

Synonyms BNSP, OPN

### **Function**

Major non-collagenous bone protein that binds tightly to hydroxyapatite. Appears to form an integral part of the mineralized matrix. Probably important to cell-matrix interaction.


**Cellular Location** Secreted.

**Tissue Location**Bone. Found in plasma.

# SPP1 Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture



Flow cytometric analysis of NCI-H292 cells using SPP1 Antibody (C-term)(bottom histogram) compared to a negative control cell (top histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

### SPP1 Antibody (C-term) - Background

SPP1 binds tightly to hydroxyapatite. The protein appears to form an integral part of the mineralized matrix and probably important to cell-matrix interaction. It acts as a cytokine involved in enhancing production of interferon-gamma and interleukin-12 and reducing production of interleukin-10 and is essential in the pathway that leads to type I immunity.

### SPP1 Antibody (C-term) - References

Macri, A., Tumori 95 (1), 48-52 (2009) Agnihotri, R., J. Biol. Chem. 276 (30), 28261-28267 (2001)