

KRT10 Antibody (N-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP6704a

Specification

KRT10 Antibody (N-term) - Product Information

| | |
|-------------------|------------------------|
| Application | WB, IHC-P, FC,E |
| Primary Accession | P13645 |
| Other Accession | P02535 |
| Reactivity | Human |
| Predicted | Mouse |
| Host | Rabbit |
| Clonality | Polyclonal |
| Isotype | Rabbit Ig |
| Calculated MW | 58827 |
| Antigen Region | 150-179 |

KRT10 Antibody (N-term) - Additional Information

Gene ID 3858

Other Names

Keratin, type I cytoskeletal 10,
Cytokeratin-10, CK-10, Keratin-10, K10,
KRT10, KPP

Target/Specificity

This KRT10 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 150-179 amino acids from the N-terminal region of human KRT10.

Dilution


WB~~1:2000
IHC-P~~1:50~100
FC~~1:10~50

Format


Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.



Anti-KRT10 Antibody (N-term) at 1:2000 dilution + HeLa whole cell lysate
Lysates/proteins at 20 µg per lane.
Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution.
Predicted band size : 59 kDa
Blocking/Dilution buffer: 5% NFDM/TBST.



Formalin-fixed and paraffin-embedded human kidney carcinoma with KRT10 Antibody (N-term), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.

Precautions

KRT10 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

KRT10 Antibody (N-term) - Protein Information

Name KRT10

Synonyms KPP

Function

Plays a role in the establishment of the epidermal barrier on plantar skin.

Cellular Location

Secreted, extracellular space. Cell surface.
Note=Localized on the surface of desquamated nasal epithelial cells (PubMed:12427098). Localized on the surface of lung cell lines (PubMed:19627498)


Tissue Location

Seen in all suprabasal cell layers including stratum corneum. Expressed on the surface of lung cell lines (PubMed:19627498).

KRT10 Antibody (N-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)



Flow cytometric analysis of 293 cells using KRT10 Antibody (N-term)(bottom histogram) compared to a negative control cell (top histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

KRT10 Antibody (N-term) - Background

KRT10 is a member of the type I (acidic) cytokeratin family, which belongs to the superfamily of intermediate filament (IF) proteins. Keratins are heteropolymeric structural proteins which form the intermediate filament. These filaments, along with actin microfilaments and microtubules, compose the cytoskeleton of epithelial cells. Mutations in its gene are associated with epidermolytic hyperkeratosis.

KRT10 Antibody (N-term) - References

- Morais,P., Eur J Dermatol 19 (4), 333-336 (2009)
Barcelos,A.C., J. Cutan. Pathol. 36 (6), 647-654 (2009)