

KRT14 Antibody (Center)

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP6713c

Specification

KRT14 Antibody (Center) - Product Information

Application WB, IHC-P, FC,E

Primary Accession <u>P02533</u>

Other Accession Q6IFU7, Q6IFX2,

Q6IFU8, Q9QWL7, Q04695, A1L595, Q6IFV1, Q61781 Human, Mouse

Reactivity Human, Mouse Predicted Rat, Bovine

Host Rabbit
Clonality Polyclonal
Isotype Rabbit Ig
Antigen Region 276-303

KRT14 Antibody (Center) - Additional Information

Gene ID 3861

Other Names

Keratin, type I cytoskeletal 14, Cytokeratin-14, CK-14, Keratin-14, K14, KRT14

Target/Specificity

This KRT14 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 276-303 amino acids from the Central region of human KRT14.

Dilution


WB~~1:2000 IHC-P~~1:50~100 FC~~1:10~50

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

Storage


Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw



Anti-KRT14 Antibody (Center) at 1:2000 dilution + A431 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution.

Predicted band size: 52 kDa

Blocking/Dilution buffer: 5% NFDM/TBST.



Formalin-fixed and paraffin-embedded human breast carcinoma with KRT14 Antibody (Center), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.



cycles.

Precautions

KRT14 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

KRT14 Antibody (Center) - Protein Information

Name KRT14

Function

The nonhelical tail domain is involved in promoting KRT5- KRT14 filaments to self-organize into large bundles and enhances the mechanical properties involved in resilience of keratin intermediate filaments in vitro.

Cellular Location

Cytoplasm. Nucleus. Note=Expressed in both as a filamentous pattern


Tissue Location

Expressed in the corneal epithelium (at protein level) (PubMed:26758872). Detected in the basal layer, lowered within the more apically located layers specifically in the stratum spinosum, stratum granulosum but is not detected in stratum corneum. Strongly expressed in the outer root sheath of anagen follicles but not in the germinative matrix, inner root sheath or hair (PubMed:9457912). Found in keratinocytes surrounding the club hair during telogen (PubMed:9457912).

KRT14 Antibody (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture



KRT14 Antibody (Center) (Cat.#AP6713c) flow cytometry analysis of MDA-MB231 cells (bottom histogram) compared to a negative control cell (top histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

KRT14 Antibody (Center) - Background

KRT14 is a member of the keratin family, the most diverse group of intermediate filaments. This protein, a type I keratin, is usually found as a heterotetramer with two keratin 5 molecules, a type II keratin. Together they form the cytoskeleton of epithelial cells.

KRT14 Antibody (Center) - References

Barcelos, A.C., J. Cutan. Pathol. 36 (6), 647-654 (2009) Romano, R.A., PLoS ONE 4 (5), E5623 (2009)