

GATA2 Antibody (C-term)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP6772b

Specification

GATA2 Antibody (C-term) - Product Information

Application	WB, FC,E
Primary Accession	P23769
Other Accession	P23824
Reactivity	Human
Predicted	Chicken
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit Ig
Calculated MW	50500
Antigen Region	380-407

GATA2 Antibody (C-term) - Additional Information

Gene ID 2624

Other Names

Endothelial transcription factor GATA-2,
GATA-binding protein 2, GATA2

Target/Specificity

This GATA2 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 380-407 amino acids from the C-terminal region of human GATA2.

Dilution

WB~~1:1000
FC~~1:10~50

Format


Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

Storage


Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

GATA2 Antibody (C-term) is for research use



Western blot analysis of GATA2 Antibody (C-term) (Cat. #AP6772b) in K562 cell line lysates (35ug/lane). GATA2 (arrow) was detected using the purified Pab.



Western blot analysis of GATA2 (arrow) using rabbit polyclonal GATA2 Antibody (C-term) (Cat. #AP6772b). 293 cell lysates (2 ug/lane) either nontransfected (Lane 1) or transiently transfected (Lane 2) with the GATA2 gene.

only and not for use in diagnostic or therapeutic procedures.

GATA2 Antibody (C-term) - Protein Information

Name GATA2

Function


Transcriptional activator which regulates endothelin-1 gene expression in endothelial cells. Binds to the consensus sequence 5'-AGATAG-3'.

Cellular Location

Nucleus.

Tissue Location

Endothelial cells.



Flow cytometric analysis of K562 cells using GATA2 Antibody (C-term)(bottom histogram) compared to a negative control cell (top histogram) FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

GATA2 Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

GATA2 Antibody (C-term) - Background

GATA2 is a member of the GATA family of zinc-finger transcription factors that are named for the consensus nucleotide sequence they bind in the promoter regions of target genes. This protein plays an essential role in regulating transcription of genes involved in the development and proliferation of hematopoietic and endocrine cell lineages.

GATA2 Antibody (C-term) - References

Pan,X., et.al., J. Biochem. 127 (1), 105-112 (2000)