

PRKACA Antibody (N-term K82)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP6822A

Specification

PRKACA Antibody (N-term K82) - Product Information

Application	WB, IHC-P, FC,E
Primary Accession	P17612
Other Accession	P68182 , P05383 , P68181 , P22694 , P68180 , P05131 , P27791 , P36887 , P05132 , P25321 , P00517 , Q9MZD9
Reactivity Predicted	Human Bovine, Hamster, Mouse, Pig, Rat, Sheep
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit Ig
Calculated MW	40590
Antigen Region	67-94

PRKACA Antibody (N-term K82) - Additional Information

Gene ID 5566

Other Names

cAMP-dependent protein kinase catalytic subunit alpha, PKA C-alpha, PRKACA, PKACA

Target/Specificity


This PRKACA antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 67-94 amino acids from the N-terminal region of human PRKACA.

Dilution


WB~~1:1000
IHC-P~~1:50~100
FC~~1:10~50

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation



Western blot analysis of PRKACA Antibody (N-term K82) (Cat. #AP6822a) in NCI-H460 cell line lysates (35ug/lane). PRKACA (arrow) was detected using the purified Pab.



Formalin-fixed and paraffin-embedded human testis tissue reacted with PRKACA Antibody (N-term K82), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.

followed by dialysis against PBS.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

PRKACA Antibody (N-term K82) is for research use only and not for use in diagnostic or therapeutic procedures.

PRKACA Antibody (N-term K82) - Protein Information

Name PRKACA

Synonyms PKACA

Function

Phosphorylates a large number of substrates in the cytoplasm and the nucleus (PubMed:<a href="http://www.uniprot.org/citations/15642694"

target="_blank">15642694 ,

PubMed:<a href="http://www.uniprot.org/citations/15905176"

target="_blank">15905176 ,

PubMed:<a href="http://www.uniprot.org/citations/16387847"

target="_blank">16387847 ,

PubMed:<a href="http://www.uniprot.org/citations/17333334"

target="_blank">17333334 ,

PubMed:<a href="http://www.uniprot.org/citations/17565987"

target="_blank">17565987 ,

PubMed:<a href="http://www.uniprot.org/citations/17693412"

target="_blank">17693412 ,

PubMed:<a href="http://www.uniprot.org/citations/18836454"

target="_blank">18836454 ,

PubMed:<a href="http://www.uniprot.org/citations/19949837"

target="_blank">19949837 ,

PubMed:<a href="http://www.uniprot.org/citations/20356841"


target="_blank">20356841 ,

PubMed:<a href="http://www.uniprot.org/citations/21085490"

target="_blank">21085490 ,

PubMed:<a href="http://www.uniprot.org/citations/21514275"

target="_blank">21514275 ,



PRKACA Antibody (N-term K82) (Cat. #AP6822a) flow cytometric analysis of NCI-H460 cells (right histogram) compared to a negative control cell (left histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

PRKACA Antibody (N-term K82) - Background

PRKACA phosphorylates a large number of substrates in the cytoplasm and the nucleus.

PRKACA Antibody (N-term K82) - References

Sugiyama,H., et.al., J. Biol. Chem. 267 (35), 25256-25263 (1992)

PubMed:21812984). Phosphorylates CDC25B, ABL1, NFKB1, CLDN3, PSMC5/RPT6, PJA2, RYR2, RORA, SOX9 and VASP (PubMed:15642694, PubMed:15905176, PubMed:16387847, PubMed:17333334, PubMed:17565987, PubMed:17693412, PubMed:18836454, PubMed:19949837, PubMed:20356841, PubMed:21085490, PubMed:21514275, PubMed:21812984). Regulates the abundance of compartmentalized pools of its regulatory subunits through phosphorylation of PJA2 which binds and ubiquitinates these subunits, leading to their subsequent proteolysis (PubMed:21423175). RORA is activated by phosphorylation (PubMed:21514275). Required for glucose- mediated adipogenic differentiation increase and osteogenic differentiation inhibition from osteoblasts (PubMed:19949837).

target="_blank">19949837). Involved in chondrogenesis by mediating phosphorylation of SOX9 (By similarity). Involved in the regulation of platelets in response to thrombin and collagen; maintains circulating platelets in a resting state by phosphorylating proteins in numerous platelet inhibitory pathways when in complex with NF-kappa-B (NFKB1 and NFKB2) and I-kappa-B-alpha (NFKBIA), but thrombin and collagen disrupt these complexes and free active PRKACA stimulates platelets and leads to platelet aggregation by phosphorylating VASP (PubMed:15642694, PubMed:20356841). Prevents the antiproliferative and anti-invasive effects of alpha- difluoromethylornithine in breast cancer cells when activated (PubMed:17333334). RYR2 channel activity is potentiated by phosphorylation in presence of luminal Ca(2+), leading to reduced amplitude and increased frequency of store overload-induced Ca(2+) release (SOICR) characterized by an increased rate of Ca(2+) release and propagation velocity of spontaneous Ca(2+) waves, despite reduced wave amplitude and resting cytosolic Ca(2+) (PubMed:17693412). PSMC5/RPT6 activation by phosphorylation stimulates proteasome (PubMed:17565987). Negatively regulates tight junctions (TJs) in ovarian cancer cells via CLDN3 phosphorylation (PubMed:15905176). NFKB1 phosphorylation promotes NF-kappa-B p50-p50 DNA binding (PubMed:15642694). Involved in embryonic development by down-regulating the Hedgehog (Hh) signaling pathway that determines embryo pattern formation and morphogenesis. Prevents meiosis resumption in prophase-arrested oocytes via CDC25B inactivation

by phosphorylation (By similarity). May also regulate rapid eye movement (REM) sleep in the pedunculopontine tegmental (PPT) (By similarity). Phosphorylates APOBEC3G and AICDA (PubMed:16387847, PubMed:18836454). Phosphorylates HSF1; this phosphorylation promotes HSF1 nuclear localization and transcriptional activity upon heat shock (PubMed:21085490).

Cellular Location

Cytoplasm. Cell membrane. Nucleus. Mitochondrion. Membrane; Lipid-anchor. Note=Translocates into the nucleus (monomeric catalytic subunit). The inactive holoenzyme is found in the cytoplasm Distributed throughout the cytoplasm in meiotically incompetent oocytes. Associated to mitochondrion as meiotic competence is acquired Aggregates around the germinal vesicles (GV) at the immature GV stage oocytes (By similarity). Colocalizes with HSF1 in nuclear stress bodies (nSBs) upon heat shock (PubMed:21085490). {ECO:0000250, ECO:0000269|PubMed:21085490}

Tissue Location

Isoform 1 is ubiquitous. Isoform 2 is sperm-specific and is enriched in pachytene spermatocytes but is not detected in round spermatids.

PRKACA Antibody (N-term K82) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)