

**NDRG1 Antibody (N-term)**  
**Affinity Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP6935a**

**Specification**

**NDRG1 Antibody (N-term) - Product Information**

|                   |  |
|-------------------|--|
| Application       | <b>WB, IHC-P, FC,E</b>   |
| Primary Accession | <a href="#">Q92597</a>   |
| Other Accession   | <a href="#">Q6JE36</a> , <a href="#">Q62433</a> ,<br><a href="#">Q4R4Q3</a> , <a href="#">Q3SYX0</a> |
| Reactivity        | <b>Human</b>   |
| Predicted         | <b>Bovine, Monkey,<br/>Mouse, Rat</b>  |
| Host              | <b>Rabbit</b>  |
| Clonality         | <b>Polyclonal</b>  |
| Isotype           | <b>Rabbit Ig</b>   |
| Calculated MW     | <b>42835</b>   |
| Antigen Region    | <b>12-40</b>   |

**NDRG1 Antibody (N-term) - Additional Information**

**Gene ID** 10397

**Other Names**

Protein NDRG1, Differentiation-related gene 1 protein, DRG-1, N-myc downstream-regulated gene 1 protein, Nickel-specific induction protein Cap43, Reducing agents and tunicamycin-responsive protein, RTP, Rit42, NDRG1, CAP43, DRG1, RTP

**Target/Specificity**


This NDRG1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 12-40 amino acids from the N-terminal region of human NDRG1.

**Dilution**


WB~~1:1000  
IHC-P~~1:50~100  
FC~~1:10~50

**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.



Western blot analysis of NDRG1 Antibody (N-term) (Cat. #AP6935a) in MDA-MB231 cell line lysates (35ug/lane). NDRG1 (arrow) was detected using the purified Pab.



Formalin-fixed and paraffin-embedded human prostate carcinoma with NDRG1 Antibody (N-term), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.

### Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

### Precautions

NDRG1 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

### NDRG1 Antibody (N-term) - Protein Information

**Name** NDRG1


**Synonyms** CAP43, DRG1, RTP

### Function

Stress-responsive protein involved in hormone responses, cell growth, and differentiation. Acts as a tumor suppressor in many cell types. Necessary but not sufficient for p53/TP53-mediated caspase activation and apoptosis. Has a role in cell trafficking, notably of the Schwann cell, and is necessary for the maintenance and development of the peripheral nerve myelin sheath. Required for vesicular recycling of CDH1 and TF. May also function in lipid trafficking. Protects cells from spindle disruption damage. Functions in p53/TP53-dependent mitotic spindle checkpoint. Regulates microtubule dynamics and maintains euploidy.

### Cellular Location

Cytoplasm, cytosol. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome. Nucleus. Cell membrane Note=Mainly cytoplasmic but differentially localized to other regions Associates with the plasma membrane in intestinal epithelia and lactating mammary gland. Translocated to the nucleus in a p53/TP53- dependent manner. In prostate epithelium and placental chorion, located in both the cytoplasm and in the nucleus. No nuclear localization in colon epithelium cells. In intestinal mucosa, prostate and renal cortex, located predominantly adjacent to adherens junctions Cytoplasmic with granular staining in proximal tubular cells of the kidney and salivary gland ducts. Recruits to the membrane of recycling/sorting and late endosomes via binding to phosphatidylinositol



Flow cytometric analysis of MDA-231 cells using NDRG1 Antibody (N-term)(bottom histogram) compared to a negative control cell (top histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

### NDRG1 Antibody (N-term) - Background

NDRG1 is a cytoplasmic protein involved in stress responses, hormone responses, cell growth, and differentiation. It is necessary for p53-mediated caspase activation and apoptosis.

### NDRG1 Antibody (N-term) - References

Sugiyama,N., et.al., Mol. Cell Proteomics 6 (6), 1103-1109 (2007)

4-phosphate. Associates with microtubules  
Colocalizes with TUBG1 in the centrosome.  
Cytoplasmic location increased with  
hypoxia. Phosphorylated form found  
associated with centromeres during  
S-phase of mitosis and with the plasma  
membrane

**Tissue Location**

Ubiquitous; expressed most prominently in  
placental membranes and prostate, kidney,  
small intestine, and ovary tissues Also  
expressed in heart, brain, skeletal muscle,  
lung, liver and pancreas. Low levels in  
peripheral blood leukocytes and in tissues  
of the immune system. Expressed mainly in  
epithelial cells. Also found in Schwann cells  
of peripheral neurons. Reduced expression  
in adenocarcinomas compared to normal  
tissues. In colon, prostate and placental  
membranes, the cells that border the lumen  
show the highest expression.

**NDRG1 Antibody (N-term) - Protocols**

Provided below are standard protocols that you  
may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**NDRG1 Antibody (N-term) - Citations**

- [Identification of differential phosphorylation and sub-cellular localization of the metastasis suppressor, NDRG1.](#)
- [Proteolytic cleavage and truncation of NDRG1 in human prostate cancer cells, but not normal prostate epithelial cells.](#)