

DHS Polyclonal Antibody

Catalog # AP69533

Specification

DHS Polyclonal Antibody - Product Information

Application
Primary Accession
Reactivity
Host
Clonality

WB
P49366
Human
Rabbit
Polyclonal

DHS Polyclonal Antibody - Additional Information

Gene ID 1725

Other Names

DHPS; DS; Deoxyhypusine synthase; DHS

Dilution

WB~~Western Blot: 1/500 - 1/2000. ELISA: 1/20000. Not yet tested in other applications.

Format

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Storage Conditions

-20°C


DHS Polyclonal Antibody - Protein Information


Name DHPS

Synonyms DS

Function

Catalyzes the NAD-dependent oxidative cleavage of spermidine and the subsequent transfer of the butylamine moiety of spermidine to the epsilon-amino group of a critical lysine residue of the eIF-5A precursor protein to form the intermediate deoxyhypusine residue (PubMed:30661771 This is the first step of the post-translational modification of that lysine into an unusual amino acid residue named hypusine.





DHS Polyclonal Antibody - Background

Catalyzes the NAD-dependent oxidative cleavage of spermidine and the subsequent transfer of the butylamine moiety of spermidine to the epsilon-amino group of a critical lysine residue of the eIF-5A precursor protein to form the intermediate



Tel: 858.875.1900 Fax: 858.622.0609



Hypusination is unique to mature eIF-5A factor and is essential for its function.

DHS Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- <u>Immunofluorescence</u>
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

deoxyhypusine residue. This is the first step of the post-translational modification of that lysine into an unusual amino acid residue named hypusine. Hypusination is unique to mature eIF- 5A factor and is essential for its function.