

FTO Antibody (N-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP6976a

Specification

FTO Antibody (N-term) - Product Information

Application	WB, IHC-P, FC,E
Primary Accession	O9C0B1
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit Ig
Calculated MW	58282
Antigen Region	1-30

FTO Antibody (N-term) - Additional Information

Gene ID 79068

Other Names

Alpha-ketoglutarate-dependent dioxygenase FTO, 11411-, Fat mass and obesity-associated protein, FTO, KIAA1752

Target/Specificity

This FTO antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 1-30 amino acids from the N-terminal region of human FTO.

Dilution

WB~~1:1000
IHC-P~~1:50~100
FC~~1:10~50

Format


Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage


Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

FTO Antibody (N-term) is for research use only and not for use in diagnostic or



Western blot analysis of FTO Antibody (N-term) (Cat. #AP6976a) in NCI-H460 cell line lysates (35ug/lane). FTO (arrow) was detected using the purified Pab.



Formalin-fixed and paraffin-embedded human brain tissue reacted with FTO Antibody (N-term), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.

therapeutic procedures.


FTO Antibody (N-term) - Protein Information

Name FTO

{ECO:0000303|PubMed:17496892,
ECO:0000312|HGNC:HGNC:24678}

Function

RNA demethylase that mediates oxidative demethylation of different RNA species, such as mRNAs, tRNAs and snRNAs, and acts as a regulator of fat mass, adipogenesis and energy homeostasis (PubMed:22002720, PubMed:26458103, PubMed:28002401, PubMed:30197295, PubMed:26457839, PubMed:25452335). Specifically demethylates N(6)-methyladenosine (m6A) RNA, the most prevalent internal modification of messenger RNA (mRNA) in higher eukaryotes (PubMed:22002720, PubMed:26458103, PubMed:30197295, PubMed:26457839, PubMed:25452335). M6A demethylation by FTO affects mRNA expression and stability (PubMed:30197295). Also able to demethylate m6A in U6 small nuclear



Flow cytometric analysis of NCI-H292 cells using FTO Antibody (N-term)(bottom histogram) compared to a negative control cell (top histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

FTO Antibody (N-term) - Background

The precise function of FTO remains to be determined.

FTO Antibody (N-term) - References

Scott,L.J., et.al., Science 316 (5829), 1341-1345 (2007)

RNA (snRNA) (PubMed:30197295). Mediates demethylation of N(6),2'-O-dimethyladenosine cap (m6A(m)), by demethylating the N(6)- methyladenosine at the second transcribed position of mRNAs and U6 snRNA (PubMed:28002401, PubMed:30197295). Demethylation of m6A(m) in the 5'-cap by FTO affects mRNA stability by promoting susceptibility to decapping (PubMed:28002401). Also acts as a tRNA demethylase by removing N(1)-methyladenine from various tRNAs (PubMed:30197295). Has no activity towards 1-methylguanine (PubMed:20376003). Has no detectable activity towards double-stranded DNA (PubMed:20376003). Also able to repair alkylated DNA and RNA by oxidative demethylation: demethylates single-stranded RNA containing 3-methyluracil, single- stranded DNA containing 3-methylthymine and has low demethylase activity towards single-stranded DNA containing 1-methyladenine or 3- methylcytosine (PubMed:18775698, PubMed:20376003). Ability to repair alkylated DNA and RNA is however unsure in vivo (PubMed:18775698, PubMed:20376003). Involved in the regulation of fat mass, adipogenesis and body weight, thereby contributing to the regulation of body size and body fat accumulation (PubMed:18775698).

target="_blank">18775698,
PubMed:<a href="http://www.uniprot.org/citations/20376003"
target="_blank">20376003). Involved
in the regulation of thermogenesis and the
control of adipocyte differentiation into
brown or white fat cells (PubMed:<a href="http://www.uniprot.org/citations/26287746"
target="_blank">26287746).
Regulates activity of the dopaminergic
midbrain circuitry via its ability to
demethylate m6A in mRNAs (By similarity).
Plays an oncogenic role in a number of
acute myeloid leukemias by enhancing
leukemic oncogene-mediated cell
transformation: acts by mediating m6A
demethylation of target transcripts such as
MYC, CEBPA, ASB2 and RARA, leading to
promote their expression (PubMed:<a href="http://www.uniprot.org/citations/28017614"
target="_blank">28017614,
PubMed:<a href="http://www.uniprot.org/citations/29249359"
target="_blank">29249359).

Cellular Location

Nucleus. Nucleus speckle. Cytoplasm
Note=Localizes mainly in the nucleus,
where it is able to demethylate
N(6)-methyladenosine (m6A) and
N(6),2'-O-dimethyladenosine cap (m6A(m))
in U6 small nuclear RNA (snRNA),
N(1)-methyladenine from tRNAs and
internal m6A in mRNAs
(PubMed:30197295). In the cytoplasm,
mediates demethylation of m6A and
m6A(m) in mRNAs and N(1)-methyladenine
from tRNAs (PubMed:30197295).

Tissue Location

Ubiquitously expressed, with relatively high
expression in adrenal glands and brain;
especially in hypothalamus and pituitary
(PubMed:17434869, PubMed:17496892).
Highly expressed in highly expressed in
acute myeloid leukemias (AML) with
t(11;11)(q23;23) with KMT2A/MLL1
rearrangements,
t(15;17)(q21;q21)/PML-RARA, FLT3-ITD,
and/or NPM1 mutations
(PubMed:28017614).

FTO Antibody (N-term) - Protocols

Provided below are standard protocols that you
may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)