

**SELL Antibody (C-term)**  
**Affinity Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP6990B**

**Specification**

**SELL Antibody (C-term) - Product Information**

Application	<b>WB, IHC-P, FC,E</b>
Primary Accession	<a href="#">P14151</a>
Reactivity	<b>Human</b>
Host	<b>Rabbit</b>
Clonality	<b>Polyclonal</b>
Isotype	<b>Rabbit Ig</b>
Antigen Region	<b>346-372</b>

**SELL Antibody (C-term) - Additional Information**

**Gene ID 6402**

**Other Names**

L-selectin, CD62 antigen-like family member L, Leukocyte adhesion molecule 1, LAM-1, Leukocyte surface antigen Leu-8, Leukocyte-endothelial cell adhesion molecule 1, LECAM1, Lymph node homing receptor, TQ1, gp90-MEL, CD62L, SELL, LNHR, LYAM1

**Target/Specificity**

This SELL antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 346-372 amino acids from the C-terminal region of human SELL.

**Dilution**


WB~~1:2000  
IHC-P~~1:10~50  
FC~~1:25

**Format**


Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.



All lanes : Anti-SELL Antibody (C-term) at 1:2000 dilution  
Lane 1: Raji whole cell lysate  
Lane 2: Jurkat whole cell lysate  
Lane 3: K562 whole cell lysate  
Lane 4: MCF-7 whole cell lysate  
Lane 5: HL-60 whole cell lysate  
Lysates/proteins at 20 µg per lane.  
Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution.  
Predicted band size : 42 kDa  
Blocking/Dilution buffer: 5% NFDM/TBST.



Formalin-fixed and paraffin-embedded

### Precautions

SELL Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

### SELL Antibody (C-term) - Protein Information

**Name** SELL

**Synonyms** LNHR, LYAM1

### Function

Calcium-dependent lectin that mediates cell adhesion by binding to glycoproteins on neighboring cells (PubMed:<a href="http://www.uniprot.org/citations/12403782" target="\_blank">12403782</a>, PubMed:<a href="http://www.uniprot.org/citations/28489325" target="\_blank">28489325</a>, PubMed:<a href="http://www.uniprot.org/citations/28011641" target="\_blank">28011641</a>). Mediates the adherence of lymphocytes to endothelial cells of high endothelial venules in peripheral lymph nodes. Promotes initial tethering and rolling of leukocytes in endothelia (PubMed:<a href="http://www.uniprot.org/citations/12403782" target="\_blank">12403782</a>, PubMed:<a href="http://www.uniprot.org/citations/28011641" target="\_blank">28011641</a>).

### Cellular Location

Cell membrane; Single-pass type I membrane protein

### Tissue Location


Expressed in B-cell lines and T-lymphocytes.

### SELL Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)

human lymph tissue with SELL Antibody (C-term), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.



Overlay histogram showing Jurkat cells stained with AP6990b (green line). The cells were fixed with 2% paraformaldehyde (10 min). The cells were then incubated in 2% bovine serum albumin to block non-specific protein-protein interactions followed by the antibody (AP6990b, 1:25 dilution) for 60 min at 37°C. The secondary antibody used was Goat-Anti-Rabbit IgG, DyLight® 488 Conjugated Highly Cross-Adsorbed(OH191631) at 1/200 dilution for 40 min at 37°C. Isotype control antibody (blue line) was rabbit IgG (1µg/1x10<sup>6</sup> cells) used under the same conditions. Acquisition of >10,000 events was performed.

### SELL Antibody (C-term) - Background

SELL is a cell surface component that is a member of a family of adhesion/homing receptors which play important roles in leukocyte-endothelial cell interactions. The molecule is composed of multiple domains: one homologous to lectins, one to epidermal growth factor, and two to the consensus repeat units found in C3/C4 binding proteins.

### SELL Antibody (C-term) - References

Zebrowska,A., et.al., Pol J Pathol 60 (1), 26-34 (2009)

- [Flow Cytometry](#)
- [Cell Culture](#)