

PICK1 (PRKCABP) Antibody (C-term)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP7078B

Specification

PICK1 (PRKCABP) Antibody (C-term) - Product Information

Application	WB,E
Primary Accession	Q9NRD5
Other Accession	Q9EP80 , Q62083 , Q4R7Q5 , Q2T9M1
Reactivity Predicted	Human, Mouse Bovine, Monkey, Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit Ig
Antigen Region	286-316

PICK1 (PRKCABP) Antibody (C-term) - Additional Information

Gene ID 9463

Other Names

PRKCA-binding protein, Protein interacting with C kinase 1, Protein kinase C-alpha-binding protein, PICK1, PRKCABP

Target/Specificity

This PICK1 (PRKCABP) antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 286-316 amino acids from the C-terminal region of human PICK1 (PRKCABP).

Dilution


WB~1:1000

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.



Western blot analysis of hPRKCABP-C300 (Cat.#AP7078b) in HL-60, Jurkat, K562 cell line lysates (35ug/lane). PRKCABP (arrow) was detected using the purified Pab.

PICK1 (PRKCABP) Antibody (C-term) - Background

PDZ domain, but not the AH domain, of PICK1 interacts with the C termini of the GTP-bound forms of ADP-ribosylation factor-1 (ARF1) and ARF3. The interactions with ARF5 and ARF6 are weak, suggesting that the PICK1 interaction is specific for class I ARFs and that it may regulate Golgi-to-endoplasmic reticulum vesicle transport. The PDZ domain of rat Pick1 interacts with the last 10 amino acids of the short C-terminal alternative splice variants of AMPA receptor subunits. It has thus been proposed that E-S-V/I-K-I, a sequence found in these 10 amino acids, is a novel PDZ-binding motif. PRKCA phosphorylates Pick1 efficiently but binds Pick1 in both the phosphorylated and unphosphorylated states. Consistent with a neuronal role for PICK1, the mouse homolog interacts with mouse AMPA glutamate receptors and colocalizes at excitatory synapses in the brain. Metabotropic glutamate receptor-7 (mGluR7) localizes specifically to presynaptic active zones. The extreme C-terminal 3 amino acids of mGluR7 have been

Precautions

PICK1 (PRKCABP) Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

PICK1 (PRKCABP) Antibody (C-term) - Protein Information

Name PICK1

Synonyms PRKCABP

Function

Probable adapter protein that bind to and organize the subcellular localization of a variety of membrane proteins containing some PDZ recognition sequence. Involved in the clustering of various receptors, possibly by acting at the receptor internalization level. Plays a role in synaptic plasticity by regulating the trafficking and internalization of AMPA receptors. May be regulated upon PRKCA activation. May regulate ASIC1/ASIC3 channel. Regulates actin polymerization by inhibiting the actin-nucleating activity of the Arp2/3 complex; the function is competitive with nucleation promoting factors and is linked to neuronal morphology regulation and AMPA receptor (AMPA) endocytosis. Via interaction with the Arp2/3 complex involved in regulation of synaptic plasticity of excitatory synapses and required for spine shrinkage during long-term depression (LTD). Involved in regulation of astrocyte morphology, antagonistic to Arp2/3 complex activator WASL/N-WASP function.

Cellular Location

Cytoplasm, perinuclear region. Membrane; Peripheral membrane protein. Membrane; Lipid-anchor. Cell junction, synapse, postsynaptic density. Cell junction, synapse, synaptosome. Cytoplasm, cytoskeleton. Note=Also membrane-associated, present at excitatory synapses.

Tissue Location

Ubiquitous.

shown to interact with the PDZ domain of PICK1. Immunofluorescence microscopy demonstrated that both proteins are localized at excitatory synapses in hippocampal neurons, with clustering of mGluR7 at synapses requires PICK1 C-terminal PDZ-binding residues. Mutant mGluR7 lacking the PDZ-binding residues localized diffusely along axons rather than at the synapse, suggesting a role for Pick1 as a scaffolding molecule at presynaptic sites.

PICK1 (PRKCABP) Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

PICK1 (PRKCABP) Antibody (C-term) - Citations

- [PDZ protein mediated activity-dependent LTP/LTD developmental switch at rat retinocollicular synapses.](#)