

## NOS3 Polyclonal Antibody

Catalog # AP71357

### Specification

#### NOS3 Polyclonal Antibody - Product Information

Application	<b>WB</b>
Primary Accession	<a href="#">P29474</a>
Reactivity	<b>Human, Mouse, Rat</b>
Host	<b>Rabbit</b>
Clonality	<b>Polyclonal</b>

#### NOS3 Polyclonal Antibody - Additional Information

**Gene ID** 4846

#### Other Names

NOS3; Nitric oxide synthase; endothelial; Constitutive NOS; cNOS; EC-NOS; Endothelial NOS; eNOS; NOS type III; NOSIII

#### Dilution

WB~~WB 1:500-2000, IF 1:50-300, IHC 1:50-300

#### Format

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

#### Storage Conditions

-20°C

#### NOS3 Polyclonal Antibody - Protein Information






**Name** NOS3

#### Function

Produces nitric oxide (NO) which is implicated in vascular smooth muscle relaxation through a cGMP-mediated signal transduction pathway. NO mediates vascular endothelial growth factor (VEGF)-induced angiogenesis in coronary vessels and promotes blood clotting through the activation of platelets.

#### Cellular Location

Cell membrane. Membrane, caveola. Cytoplasm, cytoskeleton. Golgi apparatus.



Note=Specifically associates with actin cytoskeleton in the G2 phase of the cell cycle; which is favored by interaction with NOSIP and results in a reduced enzymatic activity






### Tissue Location




Platelets, placenta, liver and kidney.




### NOS3 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)







**NOS3 Polyclonal Antibody - Background**

Produces nitric oxide (NO) which is implicated in vascular smooth muscle relaxation through a cGMP-mediated signal transduction pathway. NO mediates vascular endothelial growth factor (VEGF)-induced angiogenesis in coronary vessels and promotes blood clotting through the activation of platelets.