

RICTOR Antibody (C-term)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP7259b

Specification

RICTOR Antibody (C-term) - Product Information

Application	WB, IHC-P,E
Primary Accession	Q6R327
Other Accession	Q6QI06
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit Ig
Calculated MW	192218
Antigen Region	1617-1650

RICTOR Antibody (C-term) - Additional Information

Gene ID 253260

Other Names

Rapamycin-insensitive companion of mTOR,
AVO3 homolog, hAVO3, RICTOR
{ECO:0000312|EMBL:EAW559801}

Target/Specificity

This RICTOR antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 1617-1650 amino acids from the C-terminal region of human RICTOR.

Dilution

WB~~1:1000
IHC-P~~1:10~50


Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.


Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions



RICTOR Antibody (C-term) (Cat. #AP7259b) western blot analysis in mouse bladder tissue lysates (35ug/lane). This demonstrates the RICTOR antibody detected the RICTOR protein (arrow).



RICTOR Antibody (C-term) (Cat. #AP7259b) immunohistochemistry analysis in formalin fixed and paraffin embedded human kidney tissue followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of RICTOR Antibody (C-term) for immunohistochemistry. Clinical relevance has

RICTOR Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

not been evaluated.

RICTOR Antibody (C-term) - Protein Information

Name RICTOR

{ECO:0000312|EMBL:EAW55980.1}

Function

Subunit of mTORC2, which regulates cell growth and survival in response to hormonal signals. mTORC2 is activated by growth factors, but, in contrast to mTORC1, seems to be nutrient-insensitive. mTORC2 seems to function upstream of Rho GTPases to regulate the actin cytoskeleton, probably by activating one or more Rho-type guanine nucleotide exchange factors. mTORC2 promotes the serum-induced formation of stress-fibers or F-actin. mTORC2 plays a critical role in AKT1 'Ser-473' phosphorylation, which may facilitate the phosphorylation of the activation loop of AKT1 on 'Thr-308' by PDK1 which is a prerequisite for full activation. mTORC2 regulates the phosphorylation of SGK1 at 'Ser-422'. mTORC2 also modulates the phosphorylation of PRKCA on 'Ser-657'. Plays an essential role in embryonic growth and development.

RICTOR Antibody (C-term) - Background

RICTOR and MTOR (FRAP1) are components of a protein complex that integrates nutrient- and growth factor-derived signals to regulate cell growth.

RICTOR Antibody (C-term) - References

Pearce,L.R., Biochem. J. 405 (3), 513-522 (2007)
Yang,Q., Genes Dev. 20 (20), 2820-2832 (2006)
Jacinto,E., Cell 127 (1), 125-137 (2006)

RICTOR Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)