

**REA (PHB2) Antibody (C-term)**  
**Affinity Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP7270b**

**Specification**

**REA (PHB2) Antibody (C-term) - Product Information**

Application	WB, IHC-P,E
Primary Accession	<a href="#">Q99623</a>
Other Accession	<a href="#">Q5XIH7</a> , <a href="#">Q35129</a> , <a href="#">Q2HJ97</a> , <a href="#">NP_009204</a>
Reactivity	Human
Predicted	Bovine, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit Ig
Calculated MW	33296
Antigen Region	197-226

**REA (PHB2) Antibody (C-term) - Additional Information**

**Gene ID** 11331

**Other Names**

Prohibitin-2, B-cell receptor-associated protein BAP37, D-prohibitin, Repressor of estrogen receptor activity, PHB2  
{ECO:0000312|EMBL:AAH147661,  
ECO:0000312|HGNC:HGNC:30306}

**Target/Specificity**

This REA (PHB2) antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 197-226 amino acids from the C-terminal region of human REA (PHB2).


**Dilution**

WB~~1:1000  
IHC-P~~1:10~50


**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

**Storage**



Western blot analysis of anti-PHB2 Antibody (C-term) (Cat. #AP7270b) in 293 cell line lysates (35ug/lane). PHB2(arrow) was detected using the purified Pab.



All lanes : Anti-PHB2 (Human C-term) at 1:1000 dilution  
Lane 1: 293 whole cell lysate  
Lane 2: HeLa whole cell lysate  
Lysates/proteins at 20 µg per lane.  
Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution.  
Predicted band size : 33 kDa  
Blocking/Dilution buffer: 5% NFDM/TBST.

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

#### **Precautions**

REA (PHB2) Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

#### **REA (PHB2) Antibody (C-term) - Protein Information**

##### **Name** PHB2

{ECO:0000312|EMBL:AAH14766.1,  
ECO:0000312|HGNC:HGNC:30306}

##### **Function**

Protein with pleiotropic attributes mediated in a cell- compartment- and tissue-specific manner, which include the plasma membrane-associated cell signaling functions, mitochondrial chaperone, and transcriptional co-regulator of transcription factors and sex steroid hormones in the nucleus.


##### **Cellular Location**

Mitochondrion inner membrane. Cytoplasm. Nucleus. Cell membrane [Isoform 2]: Mitochondrion inner membrane

#### **REA (PHB2) Antibody (C-term) - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)



Formalin-fixed and paraffin-embedded human colon carcinoma tissue reacted with PHB2 Antibody (C-term) (Cat.#AP7270b), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.

#### **REA (PHB2) Antibody (C-term) - Background**

Acts as a mediator of transcriptional repression by nuclear hormone receptors via recruitment of histone deacetylases. Functions as an estrogen receptor (ER)-selective coregulator that potentiates the inhibitory activities of antiestrogens and represses the activity of estrogens. Competes with NCOA1 for modulation of ER transcriptional activity. Probably involved in regulating mitochondrial respiration activity and in aging.

#### **REA (PHB2) Antibody (C-term) - References**

Takata,H., Curr. Biol. 17 (15), 1356-1361 (2007)  
Kasashima,K., J. Biol. Chem. 281 (47), 36401-36410 (2006)