

TGase2 Polyclonal Antibody

Catalog # AP72808

Specification

TGase2 Polyclonal Antibody - Product Information

Application	WB
Primary Accession	P21980
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal

TGase2 Polyclonal Antibody - Additional Information

Gene ID 7052

Other Names

TGM2; Protein-glutamine gamma-glutamyltransferase 2; Tissue transglutaminase; Transglutaminase C; TG(C); TGC; TGase C; Transglutaminase H; TGase H; Transglutaminase-2; TGase-2

Dilution

WB~~Western Blot: 1/500 - 1/2000.
Immunohistochemistry: 1/100 - 1/300.
ELISA: 1/20000. Not yet tested in other applications.

Format

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Storage Conditions

-20°C




TGase2 Polyclonal Antibody - Protein Information

Name


TGM2
{ECO:0000303|PubMed:17939176,
ECO:0000312|HGNC:HGNC:11778}

Function


Calcium-dependent acyltransferase that catalyzes the formation of covalent bonds between peptide-bound glutamine and various primary amines, such as gamma-amino group of peptide-bound lysine, or mono- and polyamines, thereby producing cross-linked or aminated



proteins, respectively (PubMed:9252372, PubMed:23941696, PubMed:31991788). Involved in many biological processes, such as bone development, angiogenesis, wound healing, cellular differentiation, chromatin modification and apoptosis (PubMed:1683874, PubMed:7935379, PubMed:9252372, PubMed:27270573). Acts as a protein-glutamine gamma-glutamyltransferase by mediating the cross-linking of proteins, such as ACO2, HSPB6, FN1, HMGB1, RAP1GDS1, SLC25A4/ANT1, SPP1 and WDR54 (PubMed:23941696, PubMed:24349085, PubMed:29618516, PubMed:30458214). Under physiological conditions, the protein cross-linking activity is inhibited by GTP; inhibition is relieved by Ca(2+) in response to various stresses (PubMed:7649299, PubMed:7592956, PubMed:18092889). When secreted, catalyzes cross-linking of proteins of the extracellular matrix, such as FN1 and SPP1 resulting in the formation of scaffolds (PubMed:18092889).



itations/12506096" target="_blank">12506096). Plays a key role during apoptosis, both by (1) promoting the cross-linking of cytoskeletal proteins resulting in condensation of the cytoplasm, and by (2) mediating cross-linking proteins of the extracellular matrix, resulting in the irreversible formation of scaffolds that stabilize the integrity of the dying cells before their clearance by phagocytosis, thereby preventing the leakage of harmful intracellular components (PubMed:7935379, PubMed:9252372). In addition to protein cross-linking, can use different monoamine substrates to catalyze a vast array of protein post-translational modifications: mediates aminylation of serotonin, dopamine, noradrenaline or histamine into glutamine residues of target proteins to generate protein serotonylation, dopaminylation, noradrenalinylolation or histaminylation, respectively (PubMed:23797785, PubMed:30867594). Mediates protein serotonylation of small GTPases during activation and aggregation of platelets, leading to constitutive activation of these GTPases (By similarity). Plays a key role in chromatin organization by mediating serotonylation and dopaminylation of histone H3 (PubMed:30867594, PubMed:32273471). Catalyzes serotonylation of 'Gln-5' of histone H3 (H3Q5ser) during serotonergic neuron differentiation, thereby facilitating transcription (PubMed:30867594). Acts as a mediator of neurotransmission-independent role of nuclear dopamine in ventral tegmental area (VTA) neurons: catalyzes dopaminylation of 'Gln- 5' of histone H3 (H3Q5dop), thereby regulating relapse-related transcriptional plasticity in the reward system (PubMed:<a href="http://



TGase2 Polyclonal Antibody - Background

Catalyzes the cross-linking of proteins and the conjugation of polyamines to proteins.

[/www.uniprot.org/citations/32273471](http://www.uniprot.org/citations/32273471)"
target="_blank">32273471).
Regulates vein remodeling by mediating
serotonylation and subsequent inactivation
of ATP2A2/SERCA2 (By similarity). Also acts
as a protein deamidase by mediating the
side chain deamidation of specific
glutamine residues of proteins to glutamate
(PubMed:<a href="http://www.uniprot.org/citations/9623982"
target="_blank">9623982,
PubMed:<a href="http://www.uniprot.org/citations/20547769"
target="_blank">20547769).
Catalyzes specific deamidation of protein
gliadin, a component of wheat gluten in the
diet (PubMed:<a href="http://www.uniprot.org/citations/9623982"
target="_blank">9623982). May also
act as an isopeptidase cleaving the
previously formed cross-links (PubMed:26250429,
PubMed:<a href="http://www.uniprot.org/citations/27131890"
target="_blank">27131890). Also able
to participate in signaling pathways
independently of its acyltransferase
activity: acts as a signal transducer in
alpha-1 adrenergic receptor-mediated
stimulation of phospholipase C-delta (PLCD)
activity and is required for coupling alpha-1
adrenergic agonists to the stimulation of
phosphoinositide lipid metabolism
(PubMed:<a href="http://www.uniprot.org/citations/8943303"
target="_blank">8943303).

Cellular Location

Cytoplasm, cytosol. Nucleus. Chromosome.
Secreted, extracellular space, extracellular
matrix. Cell membrane
{ECO:0000250|UniProtKB:Q9WVJ6}.
Mitochondrion. Note=Mainly localizes to the
cytosol (PubMed:9575137). Present at much
lower level in the nucleus and chromatin
(PubMed:9575137). Also secreted via a
non-classical secretion pathway to the
extracellular matrix (PubMed:27270573)

TGase2 Polyclonal Antibody - Protocols

Provided below are standard protocols that you
may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)