

MEN1 Antibody (T594)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP7415b

Specification

MEN1 Antibody (T594) - Product Information

Application	IF, WB, IHC-P,E
Primary Accession	O00255
Other Accession	Q9WVR8 , O88559 , Q0P5I0 , NP_000235
Reactivity	Human
Predicted	Bovine, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit Ig
Calculated MW	68023
Antigen Region	584-615

MEN1 Antibody (T594) - Additional Information

Gene ID 4221

Other Names
Menin, MEN1, SCG2


Target/Specificity
This MEN1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 584-615 amino acids from human MEN1.

Dilution
IF~~1:10~50
WB~~1:1000
IHC-P~~1:10~50


Format
Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

Storage
Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions



Confocal immunofluorescent analysis of MEN1 Antibody (T594)(Cat#AP7415b) with Hela cell followed by Alexa Fluor 488-conjugated goat anti-rabbit IgG (green).DAPI was used to stain the cell nuclear (blue).



Western blot analysis of anti-MEN1 Antibody (T594) (Cat.#AP7415b) in Ramos cell line lysates (35ug/lane). MEN1(arrow) was detected using the purified Pab.

MEN1 Antibody (T594) is for research use only and not for use in diagnostic or therapeutic procedures.

MEN1 Antibody (T594) - Protein Information

Name MEN1

Synonyms SCG2

Function

Essential component of a MLL/SET1 histone methyltransferase (HMT) complex, a complex that specifically methylates 'Lys-4' of histone H3 (H3K4). Functions as a transcriptional regulator. Binds to the TERT promoter and represses telomerase expression. Plays a role in TGFB1-mediated inhibition of cell-proliferation, possibly regulating SMAD3 transcriptional activity. Represses JUND-mediated transcriptional activation on AP1 sites, as well as that mediated by NFKB subunit RELA. Positively regulates HOXC8 and HOXC6 gene expression. May be involved in normal hematopoiesis through the activation of HOXA9 expression (By similarity). May be involved in DNA repair.

Cellular Location

Nucleus. Note=Concentrated in nuclear body-like structures. Relocates to the nuclear matrix upon gamma irradiation


Tissue Location

Ubiquitous.

MEN1 Antibody (T594) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)



Formalin-fixed and paraffin-embedded human prostate carcinoma tissue reacted with MEN1 Antibody (T594) (Cat.#AP7415b), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.

MEN1 Antibody (T594) - Background

Menin (MEN1) is a putative tumor suppressor associated with a syndrome known as multiple endocrine neoplasia type 1. In vitro studies have shown menin is localized to the nucleus, possesses two functional nuclear localization signals, and inhibits transcriptional activation by JunD, however, the function of this protein is not known.

MEN1 Antibody (T594) - References

Hashimoto,M., Int. J. Oncol. 33 (2), 333-340 (2008)
Vidal,A., J Eur Acad Dermatol Venereol 22 (7), 835-838 (2008)
Pieterman,C.R., Clin. Endocrinol. (Oxf) (2008)