

MIG Polyclonal Antibody
Catalog # AP74191**Specification****MIG Polyclonal Antibody - Product Information**

Application	IHC
Primary Accession	Q07325
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal

MIG Polyclonal Antibody - Additional Information**Gene ID** 4283**Other Names**

C-X-C motif chemokine 9
(Gamma-interferon-induced monokine)
(Monokine induced by interferon-gamma)
(HuMIG) (MIG) (Small-inducible cytokine B9)

Dilution

IHC~~IHC-p 1:50-200, ELISA
1:10000-20000

Format

Liquid in PBS containing 50% glycerol, 0.5%
BSA and 0.02% sodium azide.

Storage Conditions




-20°C

MIG Polyclonal Antibody - Protein Information**Name** CXCL9**Synonyms** CMK, MIG, SCYB9**Function**

Cytokine that affects the growth,
movement, or activation state of cells that
participate in immune and inflammatory
response. Chemotactic for activated T-cells.
Binds to CXCR3.

Cellular Location


Secreted.



MIG Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)



MIG Polyclonal Antibody - Background

Cytokine that affects the growth, movement, or activation state of cells that participate in immune and inflammatory response. Chemotactic for activated T-cells. Binds to CXCR3.