

p21 (CDKN1A) Antibody (C-term)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP7527b

Specification

p21 (CDKN1A) Antibody (C-term) - Product Information

Application	IF, WB, IHC-P, FC, E
Primary Accession	P38936
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit Ig
Calculated MW	18119
Antigen Region	134-164

p21 (CDKN1A) Antibody (C-term) - Additional Information

Gene ID 1026

Other Names

Cyclin-dependent kinase inhibitor 1, CDK-interacting protein 1, Melanoma differentiation-associated protein 6, MDA-6, p21, CDKN1A, CAP20, CDKN1, CIP1, MDA6, PIC1, SDI1, WAF1

Target/Specificity

This p21 (CDKN1A) antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 134-164 amino acids from the C-terminal region of human p21 (CDKN1A).

Dilution


IF~~1:10~50
WB~~1:1000
IHC-P~~1:50~100
FC~~1:10~50

Format


Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

Storage

Maintain refrigerated at 2-8°C for up to 2



Confocal immunofluorescent analysis of p21 (CDKN1A) Antibody (C-term)(Cat#AP7527b) with 293 cell followed by Alexa Fluor 488-conjugated goat anti-rabbit IgG (green). Actin filaments have been labeled with Alexa Fluor 555 phalloidin (red). DAPI was used to stain the cell nuclear (blue).



The anti-CDKN1A Pab (Cat. #AP7527b) is used in Western blot to detect CDKN1A in T-47D cell lysate.

weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

p21 (CDKN1A) Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

p21 (CDKN1A) Antibody (C-term) - Protein Information

Name CDKN1A

Synonyms CAP20, CDKN1, CIP1, MDA6, PIC1, SDI1, WA

Function

May be involved in p53/TP53 mediated inhibition of cellular proliferation in response to DNA damage. Binds to and inhibits cyclin- dependent kinase activity, preventing phosphorylation of critical cyclin-dependent kinase substrates and blocking cell cycle progression. Functions in the nuclear localization and assembly of cyclin D-CDK4 complex and promotes its kinase activity towards RB1. At higher stoichiometric ratios, inhibits the kinase activity of the cyclin D- CDK4 complex. Inhibits DNA synthesis by DNA polymerase delta by competing with POLD3 for PCNA binding (PubMed:11595739). Plays an important role in controlling cell cycle progression and DNA damage- induced G2 arrest (PubMed:9106657).

Cellular Location


Cytoplasm. Nucleus

Tissue Location


Expressed in all adult tissues, with 5-fold lower levels observed in the brain

p21 (CDKN1A) Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.



Formalin-fixed and paraffin-embedded human cancer tissue reacted with the primary antibody, which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated. BC = breast carcinoma; HC = hepatocarcinoma.



p21 (CDKN1A) Antibody (C-term) (Cat. #AP7527b) flow cytometric analysis of 293 cells (right histogram) compared to a negative control cell (left histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

p21 (CDKN1A) Antibody (C-term) - Background

CDKN1A is a potent cyclin-dependent kinase inhibitor. It binds to and inhibits the activity of cyclin-CDK2 or -CDK4 complexes, and thus functions as a regulator of cell cycle

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

progression at G1. Expression is tightly controlled by the tumor suppressor protein p53, through which this protein mediates the p53-dependent cell cycle G1 phase arrest in response to a variety of stress stimuli. This protein can interact with proliferating cell nuclear antigen (PCNA), a DNA polymerase accessory factor, and plays a regulatory role in S phase DNA replication and DNA damage repair. This protein was reported to be specifically cleaved by CASP3-like caspases, which thus leads to a dramatic activation of CDK2, and may be instrumental in the execution of apoptosis following caspase activation.

p21 (CDKN1A) Antibody (C-term) - References

Fukuchi, K., et al., Biochim. Biophys. Acta 1642(3):163-171 (2003).
Frouin, I., et al., J. Biol. Chem. 278(41):39265-39268 (2003).
Dupont, J., et al., J. Biol. Chem. 278(39):37256-37264 (2003).
Di Padova, M., et al., J. Biol. Chem. 278(38):36496-36504 (2003).
Bai, Y.Q., et al., Oncogene 22(39):7942-7949 (2003).