

FGFR4 Antibody

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP7639d

Specification

FGFR4 Antibody - Product Information

Application	WB,E
Primary Accession	P22455
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit Ig
Calculated MW	87954

FGFR4 Antibody - Additional Information

Gene ID 2264

Other Names

Fibroblast growth factor receptor 4, FGFR-4, CD334, FGFR4, JTK2, TKF

Target/Specificity

This FGFR4 antibody is generated from rabbits immunized with recombinant human FGFR4 protein.

Dilution

WB~1:1000

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.


Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

FGFR4 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

FGFR4 Antibody - Protein Information



Western blot analysis of FGFR4 (arrow) using rabbit polyclonal FGFR4 Antibody (Cat.#AP7639d). 293 cell lysates (2 ug/lane) either nontransfected (Lane 1) or transiently transfected with the FGFR4 gene (Lane 2) (Origene Technologies).

FGFR4 Antibody - Background

FGFR4 is a member of the fibroblast growth factor receptor family, where amino acid sequence is highly conserved between members and throughout evolution. FGFR family members differ from one another in their ligand affinities and tissue distribution. A full-length representative protein would consist of an extracellular region, composed of three immunoglobulin-like domains, a single hydrophobic membrane-spanning segment and a cytoplasmic tyrosine kinase domain. The extracellular portion of the protein interacts with fibroblast growth factors, setting in motion a cascade of downstream signals, ultimately influencing mitogenesis and differentiation. Although alternative splicing has been observed, there is no evidence that the C-terminal half of the IgIII domain of this protein varies between three alternate forms,

Name FGFR4**Synonyms** JTK2, TKF**Function**

Tyrosine-protein kinase that acts as cell-surface receptor for fibroblast growth factors and plays a role in the regulation of cell proliferation, differentiation and migration, and in regulation of lipid metabolism, bile acid biosynthesis, glucose uptake, vitamin D metabolism and phosphate homeostasis. Required for normal down- regulation of the expression of CYP7A1, the rate-limiting enzyme in bile acid synthesis, in response to FGF19. Phosphorylates PLCG1 and FRS2. Ligand binding leads to the activation of several signaling cascades. Activation of PLCG1 leads to the production of the cellular signaling molecules diacylglycerol and inositol 1,4,5-trisphosphate. Phosphorylation of FRS2 triggers recruitment of GRB2, GAB1, PIK3R1 and SOS1, and mediates activation of RAS, MAPK1/ERK2, MAPK3/ERK1 and the MAP kinase signaling pathway, as well as of the AKT1 signaling pathway. Promotes SRC-dependent phosphorylation of the matrix protease MMP14 and its lysosomal degradation. FGFR4 signaling is down-regulated by receptor internalization and degradation; MMP14 promotes internalization and degradation of FGFR4. Mutations that lead to constitutive kinase activation or impair normal FGFR4 inactivation lead to aberrant signaling.

Cellular Location

Cell membrane; Single-pass type I membrane protein. Endosome. Endoplasmic reticulum. Note=Internalized from the cell membrane to recycling endosomes, and from there back to the cell membrane [Isoform 3]: Cytoplasm

Tissue Location

Expressed in gastrointestinal epithelial cells, pancreas, and gastric and pancreatic cancer cell lines

as indicated for family members 1-3. This particular family member preferentially binds acidic fibroblast growth factor and, although its specific function is unknown, it is overexpressed in gynecological tumor samples, suggesting a role in breast and ovarian tumorigenesis.

FGFR4 Antibody - References

Jezequel, P., et al., Br. J. Cancer 90(1):189-193 (2004).
Qian, Z.R., et al., J. Clin. Endocrinol. Metab. 89(4):1904-1911 (2004).
Yu, S., et al., J. Biol. Chem. 278(22):19597-19602 (2003).
Becker, N., et al., Cancer Epidemiol. Biomarkers Prev. 12(6):582-583 (2003).
Takenaka, H., et al., Arch. Dermatol. Res. 294(7):331-338 (2002).

FGFR4 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)