

LTK Antibody (N-term)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP7658a

Specification

LTK Antibody (N-term) - Product Information

Application	WB,E
Primary Accession	P29376
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit Ig
Antigen Region	42-72

LTK Antibody (N-term) - Additional Information

Gene ID 4058

Other Names

Leukocyte tyrosine kinase receptor, Protein tyrosine kinase 1, LTK, TYK1

Target/Specificity

This LTK antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 42-72 amino acids from the N-terminal region of human LTK.

Dilution

WB~1:2000

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.


Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

LTK Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

LTK Antibody (N-term) - Protein Information



Anti-LTK Antibody (S57) at 1:2000 dilution + human skeletal muscle lysate
Lysates/proteins at 20 µg per lane.
Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution.
Predicted band size : 92 kDa
Blocking/Dilution buffer: 5% NFDM/TBST.

LTK Antibody (N-term) - Background

Protein kinases are enzymes that transfer a phosphate group from a phosphate donor, generally the γ phosphate of ATP, onto an acceptor amino acid in a substrate protein. By this basic mechanism, protein kinases mediate most of the signal transduction in eukaryotic cells, regulating cellular metabolism, transcription, cell cycle progression, cytoskeletal rearrangement and cell movement, apoptosis, and differentiation. With more than 500 gene products, the protein kinase family is one of the largest families of proteins in eukaryotes. The family has been classified in 8 major groups based on sequence comparison of their tyrosine (PTK) or serine/threonine (STK) kinase catalytic domains. The tyrosine kinase (TK) group is mainly involved in the regulation of cell-cell

Name LTK**Synonyms** TYK1**Function**

Receptor with a tyrosine-protein kinase activity. The exact function of this protein is not known. Studies with chimeric proteins (replacing its extracellular region with that of several known growth factor receptors, such as EGFR and CSFIR) demonstrate its ability to promote growth and specifically neurite outgrowth, and cell survival. Signaling appears to involve the PI3 kinase pathway. Involved in regulation of the secretory pathway involving endoplasmic reticulum (ER) export sites (ERESs) and ER to Golgi transport.

Cellular Location

Membrane; Single-pass type I membrane protein.

Tissue Location

Expressed in non-hematopoietic cell lines and T- and B-cell lines.

interactions such as differentiation, adhesion, motility and death. There are currently about 90 TK genes sequenced, 58 are of receptor protein TK (e.g. EGFR, EPH, FGFR, PDGFR, TRK, and VEGFR families), and 32 of cytosolic TK (e.g. ABL, FAK, JAK, and SRC families).

LTK Antibody (N-term) - References

Toyoshima, H., et al., Proc. Natl. Acad. Sci. U.S.A. 90(12):5404-5408 (1993).
Krolewski, J.J., et al., EMBO J. 10(10):2911-2919 (1991).
Maru, Y., et al., Oncogene Res. 5(3):199-204 (1990).

LTK Antibody (N-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)