

**NTF3 Antibody (C-term)**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP7763b**

**Specification**

**NTF3 Antibody (C-term) - Product Information**

Application	<b>WB, FC,E</b>
Primary Accession	<a href="#">P20783</a>
Other Accession	<a href="#">P25435</a> , <a href="#">P18280</a> , <a href="#">Q06AV0</a> , <a href="#">P20181</a> , <a href="#">P25433</a> , <a href="#">Q08DT3</a>
Reactivity	<b>Human, Mouse, Rat</b>
Predicted	<b>Bovine, Chicken, Pig, Xenopus</b>
Host	<b>Rabbit</b>
Clonality	<b>Polyclonal</b>
Isotype	<b>Rabbit Ig</b>
Antigen Region	<b>152-184</b>

**NTF3 Antibody (C-term) - Additional Information**

**Gene ID** 4908

**Other Names**

Neurotrophin-3, NT-3, HDNF, Nerve growth factor 2, NGF-2, Neurotrophic factor, NTF3

**Target/Specificity**

This NTF3 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 152-184 amino acids from the C-terminal region of human NTF3.

**Dilution**

WB~~1:2000


FC~~1:25

**Format**


Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.



All lanes : Anti-NTF3 Antibody (C-term) at 1:2000 dilution  
Lane 1: Human ovary lysate  
Lane 2: Human spleen lysate  
Lysates/proteins at 20 µg per lane.  
Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution.  
Predicted band size : 29 kDa  
Blocking/Dilution buffer: 5% NFDM/TBST.



All lanes : Anti-NTF3 Antibody (C-term) at 1:2000 dilution  
Lane 1: Human brain lysate  
Lane 2: Rat brain lysate  
Lysates/proteins at

### Precautions

NTF3 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

### NTF3 Antibody (C-term) - Protein Information

**Name** NTF3

#### Function

Seems to promote the survival of visceral and proprioceptive sensory neurons.

#### Cellular Location

Secreted.

#### Tissue Location


Brain and peripheral tissues.

### NTF3 Antibody (C-term) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 29 kDa  
Blocking/Dilution buffer: 5% NFDM/TBST.



All lanes : Anti-NTF3 Antibody (C-term) at 1:2000 dilution  
Lane 1: Human liver lysate  
Lane 2: Human ovary lysate  
Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 29 kDa  
Blocking/Dilution buffer: 5% NFDM/TBST.



Anti-NTF3 Antibody (C-term) at 1:2000 dilution + NTF3 protein at 20 ng per lane.  
Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution.  
Predicted band size : 13.5 kDa  
Blocking/Dilution buffer: 5% NFDM/TBST.



Overlay histogram showing Hela cells stained with AP7763B (green line). The cells were fixed with 2% paraformaldehyde (10 min) and then permeabilized with 90% methanol for 10 min. The cells were then incubated in 2% bovine serum albumin to block non-specific protein-protein interactions followed by the antibody (AP7763B, 1:25 dilution) for 60 min at 37°C. The secondary antibody used was Goat-Anti-Rabbit IgG, DyLight® 488 Conjugated Highly Cross-Adsorbed (1583138) at 1/200 dilution for 40 min at 37°C. Isotype control antibody (blue line) was rabbit IgG1 (1 µg/1x10<sup>6</sup> cells) used under the same conditions. Acquisition of >10,000 events was performed.

### NTF3 Antibody (C-term) - Background

NTF3 is a member of the neurotrophin family that controls survival and differentiation of mammalian neurons. This protein is closely related to both nerve growth factor and brain-derived neurotrophic factor. It may be involved in the maintenance of the adult nervous system, and may affect development of neurons in the embryo when it is expressed in human placenta. NTF3-deficient mice generated by gene targeting display severe movement defects of the limbs. The mature peptide of this protein is identical in all mammals examined including human, pig, rat and mouse.

### NTF3 Antibody (C-term) - References

Hossain, W.A., J. Neurosci. Res. 86 (11), 2376-2391 (2008)  
 Mercader, J.M., Hum. Mol. Genet. 17 (9),

1234-1244 (2008)  
Kalcheim,C.,Proc. Natl. Acad. Sci. U.S.A. 89 (5),  
1661-1665 (1992)