

Methionyl tRNA Synthetase Antibody (C-term)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP7841b

Specification

**Methionyl tRNA Synthetase Antibody (C-term) -
Product Information**

Application	WB, IHC-P,E
Primary Accession	Q96GW9
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit Ig
Calculated MW	66591
Antigen Region	564-593

**Methionyl tRNA Synthetase Antibody (C-term) -
Additional Information**

Gene ID 92935

Other Names

Methionine--tRNA ligase, mitochondrial,
Methionyl-tRNA synthetase 2, Mitochondrial
methionyl-tRNA synthetase, MtMetRS,
MARS2

Target/Specificity

This Methionyl tRNA Synthetase antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 564-593 amino acids from the C-terminal region of human Methionyl tRNA Synthetase.

Dilution


WB~~1:1000
IHC-P~~1:10~50

Format


Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.



Western blot analysis of anti-MARS2 Antibody (C-term) (AP7841b) in mouse testis tissue lysates (35ug/lane). MARS2(arrow) was detected using the purified Pab.



Anti-MARS2 Antibody (C-term) at 1:1000 dilution + HL-60 whole cell lysate
Lysates/proteins at 20 µg per lane.
Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution.
Predicted band size : 67 kDa
Blocking/Dilution buffer: 5% NFDM/TBST.

Precautions

Methionyl tRNA Synthetase Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Methionyl tRNA Synthetase Antibody (C-term) - Protein Information

Name MARS2


Cellular Location

Mitochondrion matrix.

Methionyl tRNA Synthetase Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)



Formalin-fixed and paraffin-embedded human colon carcinoma tissue reacted with MARS2 antibody (C-term) (Cat.#AP7841b), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.

Methionyl tRNA Synthetase Antibody (C-term) - Background

Methionine-tRNA ligase (EC 6.1.1.10) catalyzes the chemical reaction:

ATP + L-methionine + tRNA^{Met} AMP + diphosphate + L-methionyl-tRNA^{Met}

The 3 substrates of this enzyme are ATP, L-methionine, and tRNA(Met), whereas its 3 products are AMP, diphosphate, and L-methionyl-tRNA(Met).

This enzyme participates in 3 metabolic pathways: methionine metabolism, selenoamino acid metabolism, and aminoacyl-trna biosynthesis.

Methionyl tRNA Synthetase Antibody (C-term) - References

Spencer,A.C., Biochemistry 43 (30), 9743-9754 (2004)