

**PIK3R1 Antibody (Y580)**  
**Affinity Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP8023f**

**Specification**

**PIK3R1 Antibody (Y580) - Product Information**

Application	<b>WB, IHC-P,E</b>
Primary Accession	<a href="#">P27986</a>
Other Accession	<a href="#">Q63787</a> , <a href="#">P26450</a> , <a href="#">P23727</a> , <a href="#">Q8UUU2</a>
Reactivity	<b>Human</b>
Predicted	<b>Xenopus, Bovine, Mouse, Rat</b>
Host	<b>Rabbit</b>
Clonality	<b>Polyclonal</b>
Isotype	<b>Rabbit Ig</b>
Antigen Region	<b>558-587</b>

**PIK3R1 Antibody (Y580) - Additional Information**

**Gene ID 5295**

**Other Names**

Phosphatidylinositol 3-kinase regulatory subunit alpha, PI3-kinase regulatory subunit alpha, PI3K regulatory subunit alpha, PtdIns-3-kinase regulatory subunit alpha, Phosphatidylinositol 3-kinase 85 kDa regulatory subunit alpha, PI3-kinase subunit p85-alpha, PtdIns-3-kinase regulatory subunit p85-alpha, PIK3R1, GRB1

**Target/Specificity**

This PIK3R1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 558-587 amino acids from human PIK3R1.

**Dilution**


WB~~1:1000  
IHC-P~~1:10~50

**Format**


Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

**Storage**

Maintain refrigerated at 2-8°C for up to 2



Western blot analysis of PIK3R1 (arrow) using rabbit polyclonal PIK3R1 Antibody (Y580) (Cat.#AP8023f). 293 cell lysates (2 ug/lane) either nontransfected (Lane 1) or transiently transfected with the PIK3R1 gene (Lane 2) (Origene Technologies).



Formalin-fixed and paraffin-embedded human skeletal muscle tissue reacted with PIK3R1-pY580, which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.

**PIK3R1 Antibody (Y580) - Background**

weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

#### **Precautions**

PIK3R1 Antibody (Y580) is for research use only and not for use in diagnostic or therapeutic procedures.

#### **PIK3R1 Antibody (Y580) - Protein Information**

**Name** PIK3R1

**Synonyms** GRB1

#### **Function**

Binds to activated (phosphorylated) protein-Tyr kinases, through its SH2 domain, and acts as an adapter, mediating the association of the p110 catalytic unit to the plasma membrane. Necessary for the insulin-stimulated increase in glucose uptake and glycogen synthesis in insulin-sensitive tissues. Plays an important role in signaling in response to FGFR1, FGFR2, FGFR3, FGFR4, KITLG/SCF, KIT, PDGFRA and PDGFRB. Likewise, plays a role in ITGB2 signaling (PubMed:<a href="http://www.uniprot.org/citations/17626883" target="\_blank">17626883</a>, PubMed:<a href="http://www.uniprot.org/citations/19805105" target="\_blank">19805105</a>, PubMed:<a href="http://www.uniprot.org/citations/7518429" target="\_blank">7518429</a>). Modulates the cellular response to ER stress by promoting nuclear translocation of XBP1 isoform 2 in a ER stress- and/or insulin-dependent manner during metabolic overloading in the liver and hence plays a role in glucose tolerance improvement (PubMed:<a href="http://www.uniprot.org/citations/20348923" target="\_blank">20348923</a>).

#### **Tissue Location**

Isoform 2 is expressed in skeletal muscle and brain, and at lower levels in kidney and cardiac muscle. Isoform 2 and isoform 4 are present in skeletal muscle (at protein level)

Phosphatidylinositol 3-kinase phosphorylates the inositol ring of phosphatidylinositol at the 3-prime position. The enzyme comprises a 110 kD catalytic subunit and a regulatory subunit of either 85, 55, or 50 kD. This gene encodes the 85 kD regulatory subunit.

Phosphatidylinositol 3-kinase plays an important role in the metabolic actions of insulin, and a mutation in this gene has been associated with insulin resistance.

#### **PIK3R1 Antibody (Y580) - References**

Kobayashi, H., et al., J. Biol. Chem. 279(8):6371-6379 (2004).  
Liu, H., et al., J. Cell Biol. 164(4):603-612 (2004).  
Sun, M., et al., J. Biol. Chem. 278(44):42992-43000 (2003).  
Khan, N.A., et al., J. Neurovirol. 9(6):584-593 (2003).  
Lee, H.Y., et al., J. Biol. Chem. 278(26):23630-23638 (2003).

#### **PIK3R1 Antibody (Y580) - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)