

**WEE1 Antibody (C-term)**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP8106B**

**Specification**

**WEE1 Antibody (C-term) - Product Information**

Application	<b>WB, IHC-P,E</b>
Primary Accession	<a href="#">P30291</a>
Other Accession	<a href="#">Q63802</a> , <a href="#">P47810</a>
Reactivity	<b>Human</b>
Predicted	<b>Mouse, Rat</b>
Host	<b>Rabbit</b>
Clonality	<b>Polyclonal</b>
Isotype	<b>Rabbit Ig</b>
Antigen Region	<b>604-634</b>

**WEE1 Antibody (C-term) - Additional Information**

**Gene ID** 7465

**Other Names**

Wee1-like protein kinase, WEE1hu, Wee1A kinase, WEE1

**Target/Specificity**

This WEE1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 604-634 amino acids from the C-terminal region of human WEE1.

**Dilution**

WB~~1:1000  
IHC-P~~1:50~100

**Format**


Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

**Storage**


Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

WEE1 Antibody (C-term) is for research use only and not for use in diagnostic or



Anti-WEE1 Antibody (A619) at 1:1000 dilution + A431 whole cell lysate  
Lysates/proteins at 20 µg per lane.  
Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution.  
Predicted band size : 72 kDa  
Blocking/Dilution buffer: 5% NFDM/TBST.



Formalin-fixed and paraffin-embedded human cancer tissue reacted with the primary antibody, which was peroxidase-conjugated to the secondary antibody, followed by AEC staining. This data demonstrates the use of this antibody for

therapeutic procedures.

immunohistochemistry; clinical relevance has not been evaluated. BC = breast carcinoma; HC = hepatocarcinoma.

#### **WEE1 Antibody (C-term) - Protein Information**

##### **Name WEE1**

##### **Function**

Acts as a negative regulator of entry into mitosis (G2 to M transition) by protecting the nucleus from cytoplasmically activated cyclin B1-complexed CDK1 before the onset of mitosis by mediating phosphorylation of CDK1 on 'Tyr-15'. Specifically phosphorylates and inactivates cyclin B1-complexed CDK1 reaching a maximum during G2 phase and a minimum as cells enter M phase. Phosphorylation of cyclin B1-CDK1 occurs exclusively on 'Tyr-15' and phosphorylation of monomeric CDK1 does not occur. Its activity increases during S and G2 phases and decreases at M phase when it is hyperphosphorylated. A correlated decrease in protein level occurs at M/G1 phase, probably due to its degradation.

##### **Cellular Location**

Nucleus.

#### **WEE1 Antibody (C-term) - Background**

WEE1 is a nuclear protein, which is a tyrosine kinase belonging to the Ser/Thr family of protein kinases. This protein catalyzes the inhibitory tyrosine phosphorylation of CDC2/cyclin B kinase, and appears to coordinate the transition between DNA replication and mitosis by protecting the nucleus from cytoplasmically activated CDC2 kinase.

#### **WEE1 Antibody (C-term) - References**

Kawasaki, H., et al., *Oncogene* 22(44):6839-6844 (2003).  
Hashimoto, O., et al., *Mol. Carcinog.* 36(4):171-182 (2003).  
Yuan, H., et al., *J. Virol.* 77(3):2063-2070 (2003).  
Masaki, T., et al., *Hepatology* 37(3):534-543 (2003).  
de Noronha, C.M., et al., *Science* 294(5544):1105-1108 (2001).

#### **WEE1 Antibody (C-term) - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)