

FGF4 Antibody (C-term)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP8149B

Specification

FGF4 Antibody (C-term) - Product Information

Application	WB, IHC-P,E
Primary Accession	P08620
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit Ig
Calculated MW	22048
Antigen Region	168-198

FGF4 Antibody (C-term) - Additional Information

Gene ID 2249

Other Names

Fibroblast growth factor 4, FGF-4, Heparin secretory-transforming protein 1, HST, HST-1, HSTF-1, Heparin-binding growth factor 4, HBGF-4, Transforming protein KS3, FGF4, HST, HSTF1, KS3

Target/Specificity

This FGF4 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 168-198 amino acids from the C-terminal region of human FGF4.

Dilution

WB~~1:1000
IHC-P~~1:50~100


Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.


Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions



Western blot analysis of anti-FGF4 Antibody (C-term) (Cat.#AP8149b) in CEM cell line lysates (35ug/lane). FGF4 (arrow) was detected using the purified Pab.



Formalin-fixed and paraffin-embedded human cancer tissue reacted with the primary antibody, which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated. BC = breast carcinoma; HC = hepatocarcinoma.

FGF4 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

FGF4 Antibody (C-term) - Protein Information

Name FGF4


Synonyms HST, HSTF1, KS3

Function

Plays an important role in the regulation of embryonic development, cell proliferation, and cell differentiation. Required for normal limb and cardiac valve development during embryogenesis.

Cellular Location

Secreted.



Formalin-fixed and paraffin-embedded human colon carcinoma tissue reacted with FGF4 Antibody (C-term) (Cat.#AP8149b), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.

FGF4 Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

FGF4 Antibody (C-term) - Background

FGF4 is a member of the fibroblast growth factor (FGF) family. FGF family members possess broad mitogenic and cell survival activities and are involved in a variety of biological processes including embryonic development, cell growth, morphogenesis, tissue repair, tumor growth and invasion. The gene for FGF4 was identified by its oncogenic transforming activity. The gene for FGF4 and FGF3, another oncogenic growth factor, are located closely on chromosome 11. Co-amplification of both genes was found in various kinds of human tumors. Studies on the mouse homolog suggested a function in bone morphogenesis and limb development through the sonic hedgehog (SHH) signaling pathway.

FGF4 Antibody (C-term) - References

- Yamamoto, H., et al., Oncogene 21(6):899-908 (2002).
Koh, K.R., et al., Leuk. Res. 26(10):933-938 (2002).
Sieuwerts, A.M., et al., Thromb. Haemost. 87(4):674-683 (2002).
Lopez-Sanchez, C., et al., Cell Tissue Res. 309(2):237-249 (2002).
Galland, F., et al., Cytogenet. Cell Genet. 60(2):114-116 (1992).