

BRSK1 Antibody (C-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP8168B

Specification

BRSK1 Antibody (C-term) - Product Information

Application	WB, IHC-P,E
Primary Accession	Q8TDC3
Other Accession	NP_115806
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit Ig
Calculated MW	85087
Antigen Region	654-684

BRSK1 Antibody (C-term) - Additional Information

Gene ID 84446

Other Names

Serine/threonine-protein kinase BRSK1,
Brain-selective kinase 1, Brain-specific
serine/threonine-protein kinase 1, BR
serine/threonine-protein kinase 1,
Serine/threonine-protein kinase SAD-B,
Synapses of Amphids Defective homolog 1,
SAD1 homolog, hSAD1, BRSK1, KIAA1811,
SAD1, SADB

Target/Specificity

This BRSK1 antibody is generated from
rabbits immunized with a KLH conjugated
synthetic peptide between 654-684 amino
acids from the C-terminal region of human
BRSK1.

Dilution


WB~~1:1000
IHC-P~~1:10~50

Format


Purified polyclonal antibody supplied in PBS
with 0.09% (W/V) sodium azide. This
antibody is purified through a protein A
column, followed by peptide affinity
purification.

Storage

Maintain refrigerated at 2-8°C for up to 2
weeks. For long term storage store at -20°C



KIAA1811 Antibody (S685) (Cat. #AP8168b)
western blot analysis in mouse liver tissue
lysates (35ug/lane).This demonstrates the
KIAA1811 antibody detected the KIAA1811
protein (arrow).



BRSK1 Antibody (C-term) (Cat.
#AP8168b)immunohistochemistry analysis in
formalin fixed and paraffin embedded human
testis tissue followed by peroxidase
conjugation of the secondary antibody and
DAB staining.This data demonstrates the use
of BRSK1 Antibody (C-term) for
immunohistochemistry. Clinical relevance has
not been evaluated.

BRSK1 Antibody (C-term) - Background

in small aliquots to prevent freeze-thaw cycles.

Precautions

BRSK1 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

BRSK1 Antibody (C-term) - Protein Information

Name BRSK1

Synonyms KIAA1811, SAD1, SADB

Function

Serine/threonine-protein kinase that plays a key role in polarization of neurons and centrosome duplication. Phosphorylates CDC25B, CDC25C, MAPT/TAU, RIMS1, TUBG1, TUBG2 and WEE1. Following phosphorylation and activation by STK11/LKB1, acts as a key regulator of polarization of cortical neurons, probably by mediating phosphorylation of microtubule-associated proteins such as MAPT/TAU at 'Thr-529' and 'Ser-579'. Also regulates neuron polarization by mediating phosphorylation of WEE1 at 'Ser-642' in postmitotic neurons, leading to down-regulate WEE1 activity in polarized neurons. In neurons, localizes to synaptic vesicles and plays a role in neurotransmitter release, possibly by phosphorylating RIMS1. Also acts as a positive regulator of centrosome duplication by mediating phosphorylation of gamma-tubulin (TUBG1 and TUBG2) at 'Ser-131', leading to translocation of gamma-tubulin and its associated proteins to the centrosome. Involved in the UV-induced DNA damage checkpoint response, probably by inhibiting CDK1 activity through phosphorylation and activation of WEE1, and inhibition of CDC25B and CDC25C.

Cellular Location

Cytoplasm. Nucleus. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome. Cell junction, synapse {ECO:0000250|UniProtKB:B2DD29}. Cell junction, synapse, presynaptic active zone {ECO:0000250|UniProtKB:B2DD29}. Cytoplasmic vesicle, secretory vesicle, synaptic vesicle {ECO:0000250|UniProtKB:B2DD29}

BRSK1 may be involved as a checkpoint kinase in the regulation of G2/M arrest in response to UV- or methyl methane sulfonate (MMS)-induced, but not IR-induced, DNA damage. This protein phosphorylates WEE1 and CDC25B in vitro and CDC25C in vitro and in vivo. BRSK1 is partitioned between cytoplasmic and nuclear locations in the absence of DNA damage, but translocates to the nucleus in response to Uv- or MMS-induced DNA damage. BRSK1 shares significant homology with the fission yeast Cdr2, a mitosis-regulatory kinase, and *Caenorhabditis elegans* SAD1, a neuronal cell polarity regulator. The BRSK1 transcript is expressed ubiquitously with the highest levels of expression in brain and testis.

BRSK1 Antibody (C-term) - References

Lu, R., et al., J. Biol. Chem. 279(30):31164-31170 (2004).
Lizcano, J.M., et al., EMBO J. 23(4):833-843 (2004).

Note=Nuclear in the absence of DNA damage. Translocated to the nucleus in response to UV- or MMS-induced DNA damage (By similarity)

Tissue Location

Widely expressed, with highest levels in brain and testis. Protein levels remain constant throughout the cell cycle

BRSK1 Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)