

PPP3CC Antibody (C-term)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP8465b

Specification

PPP3CC Antibody (C-term) - Product Information

Application	WB, IHC-P,E
Primary Accession	P48454
Other Accession	Q9BSS6
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit Ig
Calculated MW	58129
Antigen Region	482-512

PPP3CC Antibody (C-term) - Additional Information

Gene ID 5533

Other Names

Serine/threonine-protein phosphatase 2B catalytic subunit gamma isoform, CAM-PRP catalytic subunit, Calcineurin, testis-specific catalytic subunit, Calmodulin-dependent calcineurin A subunit gamma isoform, PPP3CC, CALNA3, CNA3

Target/Specificity

This PPP3CC antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 482-512 amino acids from the C-terminal region of human PPP3CC.

Dilution


WB~~1:1000
IHC-P~~1:50~100

Format


Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw



Western blot analysis of anti-PPP3CC Pab (Cat. #AP8465b) in A375cell line lysate.PPP3CC(arrow) was detected using the purified Pab.



Formalin-fixed and paraffin-embedded human cancer tissue reacted with the primary antibody, which was peroxidase-conjugated to the secondary antibody, followed by AEC staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated. BC = breast carcinoma; HC = hepatocarcinoma.

PPP3CC Antibody (C-term) - Background

Calmodulin-dependent protein phosphatase,

cycles.

Precautions

PPP3CC Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

PPP3CC Antibody (C-term) - Protein Information

Name PPP3CC

Synonyms CALNA3, CNA3

Function

Calcium-dependent, calmodulin-stimulated protein phosphatase which plays an essential role in the transduction of intracellular Ca(2+)-mediated signals. Dephosphorylates and activates transcription factor NFATC1. Dephosphorylates and inactivates transcription factor ELK1. Dephosphorylates DARPP32.

Tissue Location

Testis..

calcineurin, is involved in a wide range of biologic activities, acting as a Ca(2+)-dependent modifier of phosphorylation status. In testis, the motility of the sperm is thought to be controlled by cAMP-dependent phosphorylation and a unique form of calcineurin appears to be associated with the flagellum. The calcineurin holoenzyme is composed of catalytic and regulatory subunits of 60 and 18 kD, respectively. At least 3 genes, calcineurin A-alpha, calcineurin A-beta, and calcineurin A-gamma, have been cloned for the catalytic subunit. These genes have been identified in humans, mice, and rats, and are highly conserved between species (90 to 95% amino acid identity).

PPP3CC Antibody (C-term) - References

Eastwood, S.L., et al., Biol. Psychiatry 57(7):702-710 (2005).
Gerber, D.J., et al., Proc. Natl. Acad. Sci. U.S.A. 100(15):8993-8998 (2003).
Bennasser, Y., et al., Virology 303(1):174-180 (2002).
Esau, C., et al., J. Exp. Med. 194(10):1449-1459 (2001).
Muramatsu, T., et al., Biochem. Biophys. Res. Commun. 188(1):265-271 (1992).

PPP3CC Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)