

**ABI1 Antibody (N-term)**  
**Affinity Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP8535A**

**Specification**

**ABI1 Antibody (N-term) - Product Information**

Application	IF, WB, IHC-P, FC,E
Primary Accession	<a href="#">Q8IZP0</a>
Other Accession	<a href="#">P62484</a> , <a href="#">Q9NYB9</a> , <a href="#">Q9QZM5</a> , <a href="#">Q8CBW3</a> , <a href="#">F1R187</a>
Reactivity Predicted	Human Zebrafish, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit Ig
Calculated MW	55081
Antigen Region	81-108

**ABI1 Antibody (N-term) - Additional Information**

**Gene ID** 10006

**Other Names**

Abl interactor 1, Abelson interactor 1, Abi-1,  
Abl-binding protein 4, AblBP4, Eps8 SH3  
domain-binding protein, Eps8-binding  
protein, Nap1-binding protein, Nap1BP,  
Spectrin SH3 domain-binding protein 1,  
e3B1, ABI1, SSH3BP1

**Target/Specificity**


This ABI1 antibody is generated from  
rabbits immunized with a KLH conjugated  
synthetic peptide between 81-108 amino  
acids from the N-terminal region of human  
ABI1.

**Dilution**


IF~~1:10~50  
WB~~1:1000  
IHC-P~~1:50~100  
FC~~1:10~50

**Format**

Purified polyclonal antibody supplied in PBS  
with 0.09% (W/V) sodium azide. This  
antibody is purified through a protein A  
column, followed by peptide affinity



Immunofluorescence analysis of ABI1  
Antibody (N-term) with paraffin-embedded  
human brain tissue . 0.05 mg/ml primary  
antibody was followed by FITC-conjugated  
goat anti-rabbit IgG (whole molecule). FITC  
emits green fluorescence.



Western blot analysis of ABI1 Antibody  
(N-term) (Cat. #AP8535a) in MCF-7, CEM,  
Jurkat cell line lysates (35ug/lane). ABI1  
(arrow) was detected using the purified Pab.

purification.

### Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

### Precautions

ABI1 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

### ABI1 Antibody (N-term) - Protein Information

**Name** ABI1 ([HGNC:11320](#))

**Synonyms** SSH3BP1

### Function


May act in negative regulation of cell growth and transformation by interacting with nonreceptor tyrosine kinases ABL1 and/or ABL2. May play a role in regulation of EGF-induced Erk pathway activation. Involved in cytoskeletal reorganization and EGFR signaling. Together with EPS8 participates in transduction of signals from Ras to Rac. In vitro, a trimeric complex of ABI1, EPS8 and SOS1 exhibits Rac specific guanine nucleotide exchange factor (GEF) activity and ABI1 seems to act as an adapter in the complex. Regulates ABL1/c-Abl- mediated phosphorylation of ENAH. Recruits WASF1 to lamellipodia and there seems to regulate WASF1 protein level. In brain, seems to regulate the dendritic outgrowth and branching as well as to determine the shape and number of synaptic contacts of developing neurons.

### Cellular Location


Cytoplasm. Nucleus. Cell projection, lamellipodium. Cell projection, filopodium. Cell projection, growth cone Cell junction, synapse, postsynaptic density. Cytoplasm, cytoskeleton. Note=Localized to protruding lamellipodia and filopodia tips. Also localized to neuronal growth cones and synaptosomes. May shuttle from the postsynaptic densities to the nucleus (By similarity).

### Tissue Location

Widely expressed, with highest expression in brain.



Formalin-fixed and paraffin-embedded human brain tissue reacted with ABI1 Antibody (N-term), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.



Flow cytometric analysis of SK-Br-3 cells using ABI1 Antibody (N-term)(bottom histogram) compared to a negative control cell (top histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

### ABI1 Antibody (N-term) - Background

ABI1 has been found to form a complex with EPS8 and SOS1, and is thought to be involved in the transduction of signals from Ras to Rac. In addition, this protein may play a role in the regulation of EGF-induced Erk pathway activation as well as cytoskeletal reorganization and EGFR signaling.

### ABI1 Antibody (N-term) - References

**ABI1 Antibody (N-term) - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Wang,C., et.al., Mol. Cancer Res. 5 (10),  
1031-1039 (2007)  
Carabeo,R.A., et.al., Cell. Microbiol. 9 (9),  
2278-2288 (2007)