

ITGA1 Antibody (Center)

Affinity Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP8814C

Specification

ITGA1 Antibody (Center) - Product Information

Application	WB, IHC-P,E
Primary Accession	<u>P56199</u>
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit Ig
Calculated MW	130848
Antigen Region	727-755

ITGA1 Antibody (Center) - Additional Information

Gene ID 3672

Other Names

Integrin alpha-1, CD49 antigen-like family member A, Laminin and collagen receptor, VLA-1, CD49a, ITGA1

Target/Specificity

This ITGA1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 727-755 amino acids from the Central region of human ITGA1.

Dilution WB~~1:1000 IHC-P~~1:10~50

Format


Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage


Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

ITGA1 Antibody (Center) is for research use only and not for use in diagnostic or



Western blot analysis of ITGA1 Antibody (Center) (Cat. #AP8814c) in mouse cerebellum tissue lysates (35ug/lane). ITGA1 (arrow) was detected using the purified Pab.



Formalin-fixed and paraffin-embedded human brain tissue reacted with ITGA1 Antibody (Center), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has



therapeutic procedures.

ITGA1 Antibody (Center) - Protein Information

Name ITGA1

Function

Integrin alpha-1/beta-1 is a receptor for laminin and collagen. It recognizes the proline-hydroxylated sequence G-F-P-G-E-R in collagen. Involved in anchorage-dependent, negative regulation of EGF-stimulated cell growth.

Cellular Location Membrane; Single-pass type I membrane protein.

ITGA1 Antibody (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- <u>Blocking Peptides</u>
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

ITGA1 Antibody (Center) - Citations

• <u>Differential expression of sema3A and sema7A in a murine model of multiple sclerosis:</u> <u>Implications for a therapeutic design.</u>

not been evaluated.

ITGA1 Antibody (Center) - Background

ITGA1 is the alpha 1 subunit of integrin receptors. This protein heterodimerizes with the beta 1 subunit to form a cell-surface receptor for collagen and laminin. The heterodimeric receptor is involved in cell-cell adhesion and may play a role in inflammation and fibrosis. The alpha 1 subunit contains an inserted (I) von Willebrand factor type I domain which is thought to be involved in collagen binding.

ITGA1 Antibody (Center) - References

Bank,I., et.al., J. Clin. Immunol. 11 (1), 29-38 (1991)