

ELF3 Antibody (C-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP9441b

Specification

ELF3 Antibody (C-term) - Product Information

Application	WB,E
Primary Accession	P78545
Other Accession	Q4V7E1 , Q3UPW2 , O70273 , Q9NZC4 , Q32LN0
Reactivity	Human
Predicted	Bovine, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit Ig
Calculated MW	41454
Antigen Region	322-349

ELF3 Antibody (C-term) - Additional Information

Gene ID 1999

Other Names

ETS-related transcription factor Elf-3,
E74-like factor 3, Epithelial-restricted with
serine box, Epithelium-restricted Ets protein
ESX, Epithelium-specific Ets transcription
factor 1, ESE-1, ELF3 ([http://www.
genenames.org/cgi-bin/gene_symbol_report
?hgnc_id=3318](http://www.genenames.org/cgi-bin/gene_symbol_report?hgnc_id=3318)
target="_blank">HGNC:3318)

Target/Specificity


This ELF3 antibody is generated from
rabbits immunized with a KLH conjugated
synthetic peptide between 322-349 amino
acids from the C-terminal region of human
ELF3.

Dilution

WB~~1:1000

Format

Purified polyclonal antibody supplied in PBS
with 0.09% (W/V) sodium azide. This
antibody is purified through a protein A
column, followed by peptide affinity
purification.



Western blot analysis of ELF3 Antibody (C-term) (Cat. #AP9441b) in MCF-7 cell line lysates (35ug/lane). ELF3 (arrow) was detected using the purified Pab.

ELF3 Antibody (C-term) - References

- Lee, S.H., et al. Mol. Cancer Ther. 7(12):3739-3750(2008)
- Wu, J., et al. Cell Res. 18(6):649-663(2008)
- Peng, H., et al. J. Cell. Physiol. 215(2):562-573(2008)
- Iwai, S., et al. Oral Dis 14(2):144-149(2008)
- Manavathi, B., et al. J. Biol. Chem. 282(27):19820-19830(2007)
- Choi, S.G., et al. J. Biol. Chem. 273(1):110-117(1998)

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

ELF3 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

ELF3 Antibody (C-term) - Protein Information

Name ELF3 ([HGNC:3318](#))

Function

Transcriptional activator that binds and transactivates ETS sequences containing the consensus nucleotide core sequence GGA[AT]. Acts synergistically with POU2F3 to transactivate the SPRR2A promoter and with RUNX1 to transactivate the ANGPT1 promoter. Also transactivates collagenase, CCL20, CLND7, FLG, KRT8, NOS2, PTGS2, SPRR2B, TGFBR2 and TGM3 promoters. Represses KRT4 promoter activity. Involved in mediating vascular inflammation. May play an important role in epithelial cell differentiation and tumorigenesis. May be a critical downstream effector of the ERBB2 signaling pathway. May be associated with mammary gland development and involution. Plays an important role in the regulation of transcription with TATA-less promoters in preimplantation embryos, which is essential in preimplantation development (By similarity).

Cellular Location

Cytoplasm. Nucleus
{ECO:0000255|PROSITE-ProRule:PRU00237,
ECO:0000269|PubMed:10391676,
ECO:0000269|PubMed:15169914,
ECO:0000269|PubMed:17060315}

Note=Localizes to the cytoplasm where it has been shown to transform MCF-12A mammary epithelial cells via a novel cytoplasmic mechanism Also transiently expressed and localized to the nucleus where it induces apoptosis in non-transformed breast epithelial cells MCF-10A and MCF-12A via a transcription-dependent mechanism

Tissue Location

Expressed exclusively in tissues containing

a high content of terminally differentiated epithelial cells including mammary gland, colon, trachea, kidney, prostate, uterus, stomach and skin

ELF3 Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)