


CXCR4 Antibody

Catalog # ASC10002

Specification

CXCR4 Antibody - Product Information

Application	IF
Primary Accession	CXCR4
Other Accession	NP_003458 , 7852
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Calculated MW	39 kDa KDa
Application Notes	CXCR4 antibody can be used for Western blot 0.5 - 1 µg/mL. Antibody can also be used for Immunohistochemistry starting at 5 µg/mL. For immunofluorescence start at 4 µg/mL.



Immunofluorescence of NogoA in mouse brain tissue with NogoA Antibody at 20 µg/mL.

CXCR4 Antibody - Additional Information

Gene ID **7852**
Other Names
CXCR4 Antibody: FB22, HM89, LAP3, LCR1, NPYR, WHIM, CD184, LESTR, NPY3R, NPYRL, HSY3RR, NPYY3R, D2S201E, chemokine (C-X-C motif) receptor 4

Target/Specificity

CXCR4 antibody was raised against a 15 amino acid peptide near the center of human CXCR4. The immunogen is located within amino acids 170 - 220 of CXCR4.

Reconstitution & Storage

CXCR4 antibody can be stored at 4°C for three months and -20°C, stable for up to one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.

Precautions

CXCR4 Antibody is for research use only

CXCR4 Antibody - Background

CXCR4 Antibody: Human immunodeficiency virus (HIV) and related viruses require coreceptors, in addition to CD4, to infect target cells. Some G protein-coupled receptors including CCR5, CXCR4, CCR3, CCR2b and CCR8 in the chemokine receptor family, and four new human molecules GPR15, STRL33, GPR1 and V28 were recently identified as HIV coreceptors. Among them, CXCR4 (fusin, LESTR or HUMSTR) is a principal coreceptor for T-cell tropic strains of HIV-1 fusion and entry of human white blood cells. CXCR4 is also required for the infection by dual-tropic strains of HIV-1 and mediates CD-4 independent infection by HIV-2. The α -chemokine SDF-1 is the ligand for CXCR4 and prevents infection by T-tropic HIV-1. CXCR4 associates with the surface CD4-gp120 complex before HIV enters target cells. CXCR4 messenger RNA levels correlated with HIV-1 permissiveness in diverse human cell types. Antibodies to CXCR4 block HIV-1 and HIV-2 fusion and infection of human target cells. The amino-terminal domain and the second extracellular loop of CXCR4 serve

and not for use in diagnostic or therapeutic procedures.

as HIV binding sites.

CXCR4 Antibody - Protein Information

CXCR4 Antibody - References

Dimitrov DS. Cell 1997;91:721-730
Feng Y et al. Science 1996;272:872-7
Berson JF et al. J Virol 1996;70:6288-95
Doranz BJ et al. Cell 1996;85:1149-1158

CXCR4 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)