

TCCR Antibody

Catalog # ASC10147

Specification

TCCR Antibody - Product Information

Application	WB, IHC, IF
Primary Accession	O6UWB1
Other Accession	NP_004834 , 4759328
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Calculated MW	70 kDa KDa
Application Notes	TCCR antibody can be used for detection of TCCR by Western blot at 1 µg/mL. An approximately 70 kDa band can be detected. Antibody can also be used for immunohistochemistry starting at 5 µg/mL. For immunofluorescence start at 20 µg/mL.


TCCR Antibody - Additional Information

Gene ID **9466**
Other Names
TCCR Antibody: CRL1, TCCR, WSX1, IL27R, IL-27RA, zcytor1, CRL1, UNQ296/PRO336, Interleukin-27 receptor subunit alpha, Cytokine receptor WSX-1, IL-27 receptor subunit alpha, interleukin 27 receptor, alpha


Target/Specificity
IL27RA;

Reconstitution & Storage


TCCR antibody can be stored at 4°C for three months and -20°C, stable for up to one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.



Western blot analysis of TCCR expression in human spleen tissue lysate with TCCR antibody at 1 µg /ml.



Immunohistochemistry of TCCR in human spleen tissue with TCCR antibody at 5 µg/mL.



Immunofluorescence of TCCR in Human Spleen cells with TCCR antibody at 20 µg/mL.

TCCR Antibody - Background

TCCR Antibody: Upon antigen challenge, T-helper cells differentiate into two functional distinct subsets, Th1 and Th2. Th1 cells produce IL-2, IFN-gamma and lymphotoxin-beta that augment cell mediated

Precautions

TCCR Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

TCCR Antibody - Protein Information

Name IL27RA

Synonyms CRL1, TCCR, WSX1

Function

Receptor for IL27. Requires IL6ST/gp130 to mediate signal transduction in response to IL27. This signaling system acts through STAT3 and STAT1. Involved in the regulation of Th1-type immune responses. Also appears to be involved in innate defense mechanisms.

Cellular Location

Membrane; Single-pass type I membrane protein.

Tissue Location

Highly expressed in lymphoid tissues such as spleen, lymph nodes and peripheral blood leukocytes. Weakly expressed in other tissues examined including heart, brain, fetal and adult lung, liver, skeletal muscle, kidney, pancreas, prostate, testis, ovary, small intestine, kidney and colon. In the lymphoid system, higher level expression in CD4+ T-cell subsets than in CD8+ T-cell subsets. Also weaker expression in CD19+ B-cells and monocytes

immune response while Th2 cells secrete IL-4, IL-5, and IL-10 that enhance humoral immunity. The function of T-helper cells is regulated by cytokines. A novel cytokine receptor was recently identified and cloned. It is a new member in the type I cytokine receptor family and designated TCCR for T-cell cytokine receptor and WSX-1. TCCR deficient mice had impaired Th1 responses to protein antigen challenge, including decreased levels of IFN-gamma and Th1-dependent antibody IgG2a. TCCR is predominately expressed in thymus, spleen, lymph nodes and peripheral blood leukocytes.

TCCR Antibody - References

Chen Q, Ghilardi N, Wang H, Baker T, Xie MH, Gurney A, Grewal IS and de Sauvage FJ. Development of Th1-type immune responses requires the type I cytokine receptor TCCR Nature 2000;407(6806):916-920
Sprecher, C.A., Grant, F.J., Baumgartner, J.W., Presnell, S.R., Schrader, S.K., Yamagiwa, T., Whitmore, T.E., O'Hara, P.J. and Foster, D.F. Cloning and characterization of a novel class I cytokine receptor Biochem. Biophys. Res. Commun. 1998;246(1):82-90

TCCR Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)